

Hora	Duración	Ponencia	Título	Ponente
9:15	15'	Organización Encuentro		
9:30	15'	Sistemas Acuosos I – Clatratos: Presentación del Moderador		Olga Prieto Ballesteros
9:45	2x15'	Ponencias Sistemas Acuosos I	Clatratos en el Sistema Solar	Mª Victoria Muñoz Iglesias
			Análisis cualitativo y cuantitativo del CO ₂ en salmueras	Laura Jiménez Bonales
10:15	45'	Conferencia Invitada	Emerging risks related to food technology	Jean Claude Cheftel
11:00	15'	Sistemas Acuosos II – Disoluciones: Presentación del Moderador		Pedro D. Sanz Martínez
11:15	3x15'	Ponencias Sistemas Acuosos II	Efectos inducidos por la presión sobre agregados en medio acuoso. Estudio del dodecanoato de sodio mediante medidas de velocidad del sonido	Eduardo Hidalgo Baltasar
			Efectos inducidos por la presión sobre agregados en medio acuoso. Aplicación de técnicas volumétricas al estudio de la asociación de caseínas	Bérengère Guignon
			Influencia de la presión en la simulación del plegamiento de proteínas	Ramiro Perezzan Rodríguez
12:00	Pausa			
12:30	15'	Sistemas Acuosos III – Microbiología y T. Alimentos: Presentación		Buenaventura Guamis
12:45	4x15'	Ponencias Sistemas Acuosos III	Muerte y daño subletal causados por alta presión	Jordi Saldo Periago
			Nanoencapsulación por Ultra Alta Presión Homogenización (UHPH)	Roger Escriu Justo
			Incorporación de Sensores Ópticos al procesado de alimentos	Manuel Castillo Zambudio
			Modelización matemática de procesos a alta presión	Nadia A. S. Smith
13:45	15'	Sistemas Acuosos - Resumen, Discusión y Perspectivas		
14:15	Comida			
15:30	Sesiones 3M – Round Tables 1-8			
16:30	45	Conferencia Invitada	Magnetism in oxygen under pressure	Stefan Klotz
17:15	15'	Sistemas Moleculares I – Materiales NOCH: Presentación del Moderador		Mercedes Taravillo Corralo
17:30	3x15'	Ponencias Sistemas Moleculares I	Caracterización de yunques de zafiro y moissanita a alta presión	Elena del Corro García
			Presión interna de disoluciones acuosas	Nubia J. Mendoza Lugo
			Síntesis de Resinas Fósiles a alta presión. Resina, copal y ámbar	Óscar R. Montoro
18:15	Pausa			
18:45	15'	Sistemas Moleculares II – Enlace y Reactividad: Presentación		Víctor Luaña Cabal
19:00	4x15'	Ponencias Sistemas Moleculares II	Funciones de distribución del número de electrones (EDF)	Evelio Francisco Miguélez
			Enlace químico en moléculas heterodiatómicas bajo la perspectiva de Átomos Cuánticos Interaccionantes	Aurora Costales Castro
			Modelización de la alta presión con métodos de función de onda	José Tortajada Pérez
			Desarrollo de técnicas híbridas QM/MM y dinámica molecular para el estudio de la estructura y reactividad	José Javier Ruiz Pernía
20:00	15'	Sistemas Moleculares - Resumen, Discusión y Perspectivas		
20:15	45'	Conferencia Invitada	Nanociencia Molecular y Alta Presión	Eugenio Coronado Miralles
21:00	Cena			
22:30	Sesiones 3M – Round Tables 9-16			

Martes 25 de Enero


Hora	Duración	Ponencia	Ponente
9:30	10'	Infraestructuras MALTA-Consolider: Presentación del Moderador	Óscar R. Montoro
9:40	20'	Infraestructura I – Servidor MALTA	José Manuel Menéndez Montes
10:00	20'	Infraestructura II – Difracción de Rayos X	David Santamaría Pérez
10:20	20'	Infraestructura III – Raman Confocal	Jesús González Gómez
10:40	20'	Infraestructura IV – FTIR	Vinod Panchal
11:00	20'	Infraestructura V – Sistema de Calentamiento Láser	Daniel Errandonea
11:20	20'	Infraestructura VI – Prensa de Gran Volumen	Ramón Gervilla Fernández
11:40	20'	Infraestructura VII – Centro de Mecanizado	Samuel Gilliland
12:00	Pausa		
12:30	15'	Edición del Libro Especializado en Altas Presiones	Rafael Valiente Barroso
12:45	15'	V Escuela de Altas Presiones – La Laguna 2011	Alfonso Muñoz/Víctor Lavín
13:00	15'	Resumen Reunión Sociedad Francesa de Alta Presión	Jesús A. González Gómez
13:15	15'	Organización Congreso AIRAPT/EHPRG Madrid 2015	Fernando Rodríguez González
14:00	Comida		
15:30	Sesiones 3M –Cross-Link Round Tables A		
16:30	Sesión Monográfica: Instrumentación a Alta Presión		
16:30	30'	Homenaje a Jean-Claude Chervin	Alfredo Segura García del Río Jesús A. González Gómez
17:00	45'	Conferencia Invitada	Desarrollo de Dispositivos de Alta Presión Jean-Claude Chervin
17:45	Pausa		
18:00	20'	Celdas París-Edimburgo	Domingo Martínez García
18:20	20'	Status of the High Pressure end-station at Alba	Inmaculada Peral Alonso
18:40	20'	Desarrollo en celdas de alta presión para medidas de propiedades magnéticas	Javier Sánchez Benítez
19:00	20'	Simulación de Procesos a alta presión	Ángel Ramos del Olmo
19:20	Resumen Tercer Año MALTA-Evaluación MICINN		
20:30	Salida Cena Oficial		
21:00	Cena Oficial de la Reunión		
23:30	Regreso Cena Oficial a la Sede		

Miércoles 26 de Enero

Hora	Duración	Ponencia		Ponente
9:30	15'	Óxidos y Minerales I –Estructura y Reactividad 1: Presentación del Moderador		Alfredo Segura García del Río
9:45	4x15'	Ponencias Óxidos y Minerales I – 1	Aplicación de la absorción de rayos X al estudio de las delafositas bajo presión	Julio Pellicer Porres
			Estudio estructural de la distorsión Jahn-Teller del CuWO ₄ bajo presión	Javier Ruiz Fuertes
			Transición de fase en Rb ₂ S a alta presión. Comparación estructural de los sulfuros alcalinos y sus óxidos correspondientes	David Santamaría Pérez
			Estudio bajo altas presiones de la estructura de minerales de la corteza y de la capa superior del manto	Daniel Errandonea
10:45	45'	Conferencia Invitada	Difracción de Neutrones en Condiciones Extremas: XTREMED	Javier Campo Ruiz
11:30	15'	Óxidos y Minerales I – Estructura y Reactividad 2: Presentación del Moderador		Alfonso Muñoz González
11:30	4x15'	Ponencias Óxidos y Minerales I – 2	HgWO ₄ under pressure	Javier López Solano
			Estudio del MnGa ₂ Se ₄ bajo presión	Eduardo Pérez González
			Materiales de alta presión como electrodo para baterías de litio	Elena Arroyo y de Dompablo
12:30	Pausa			
13:00	15'	Óxidos y Minerales I – Estructura y Reactividad 3: Presentación del Moderador		José Manuel Recio Muñiz
13:15	5x12'	Ponencias Óxidos y Minerales I – 3	Rigorous treatment of first principles data for quasiharmonic thermodynamics of solids	Alberto Otero de la Roza
			Más allá del modelo de Debye. Nuevo módulo de cálculo de efectos térmicos en cristales	David Abassi Pérez
			Estudio sistemático de las ecuaciones de estado en las perovskitas KMF ₃	José Manuel Menéndez Montes
			Estudio de polimorfos de TiO ₂ . DFT+U	Ángel Morales García
			Metales alcalinos a alta presión	Miriam Marqués Arias
14:30	Comida			
16:00	Sesiones 3M –Cross-Link Round Tables B			
17:00	15'	Óxidos y Minerales I – Estructura y Reactividad 4: Presentación del Moderador		Armando Beltran Flors
17:15	4x15'	Ponencias Óxidos y Minerales I – 4	Modelización de transiciones de fase inducidas por presión en compuestos ABO ₄ y A ₂ BO ₄	Lourdes Gracia Edo
			Comportamiento estructural de perovskitas sometidas a condiciones extremas	Fernando Aguado Menéndez
			Estudio teórico (y experimental) del efecto de la presión sobre TiO ₂ y SbPO ₄	Marcilene Cristina Gomes
			Síntesis y caracterización de perovskitas complejas derivadas del CaCu ₃ Mn ₄ O ₁₂	Paula Kayser González
18:15	15'	Óxidos y Minerales I –Estructura y Reactividad – Resumen, Discusión y Perspectivas		
18:30	Pausa			
19:00	30'	Miguel Álvarez Blanco – Recuerdo/Homenaje		Ángel Martín Pendás
19:30	90'	Reunión Comités MALTA-Consolider		
21:00	Cena			
22:30	Sesiones 3M –Round Table Final			

Jueves 27 de Enero

Hora	Duración	Ponencia	Ponente	
9:30	15'	Óxidos y Minerales II – Espectroscopia 1: Presentación del Moderador		
9:45	5x15'	Ponencias Óxidos y Minerales II – Espectroscopia 1	Metastability of wurtzite and rock-salt phases in ZnCoO nanoparticles investigated by optical absorption	Rosa Martín Rodríguez
			Transiciones de spin en sistemas Jahn-Teller de Mn^{3+} , Ni^{3+} , Co^{3+}	Marta N. Sanz Ortiz
			Luminescence dependence of NaYF ₄ :Er ³⁺ , Yb ³⁺ with pressure	Carlos Renero Lecuna
			Absorción óptica bajo presión del compuesto multiferroico BiFeO ₃ . Estudio del momento dipolar eléctrico	Susana Gómez Salces
			Medidas ópticas, magnéticas y vibracionales en películas de ZnO:Co co-dopadas con Ga	Gloria Almonacid Caballer
11:00	15'	Óxidos y Minerales II – Espectroscopia 2: Presentación del Moderador		
11:15	2x15'	Ponencias Óxidos y Minerales II – Espectroscopia 2	High-pressure study of YVO ₄ nanoboxes	Óscar Gomis Hilario
			High-Pressure Raman Scattering in AB ₂ X ₄ compounds	Rosario Vilaplana Cerdá
12:00	Pausa			
11:45	15'	Óxidos y Minerales II – Espectroscopia 3: Presentación del Moderador		
12:30	4x15'	Ponencias Óxidos y Minerales II – Espectroscopia 3	Espectroscopia a alta presión en granates nanocristalinos	Ulises Rodríguez Mendoza
			Desarrollo de nuevos materiales prácticamente activos para sensores de presión	Sergio F. León Luis
			Seguimiento de cambios estructurales bajo presión mediante campo cristalino	Juan E. Muñoz Santiuste
			Desarrollo de materiales ópticamente activos como sensores de temperatura	Inocencio R. Martín Benenzuela
13:30	15'	Óxidos y Minerales II – Espectroscopia - Resumen, Discusión y Perspectivas		
13:45	Entrega Certificados Simposio y Sesiones 3M			
	Clausura del Simposio			
14:00	Comida			
16:00	Salida Autobús			



SISTEMAS ACUOSOS –I

AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

www.malta-consolider.com



Matter at High Pressure



Grupo: Centro de Astrobiología

Felipe Gómez
Laura Jiménez Bonales
José Antonio R. Manfredi
Victoria Muñoz Iglesias
Olga Prieto Ballesteros

I. AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

a. Síntesis, propiedades y estabilidad de clatratos

b. Efectos de la alta presión en soluciones acuosas de agregados supramoleculares y proteínas

c. Microbiología en condiciones extremas de presión y temperatura

I. AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

a. Síntesis, propiedades y estabilidad de clatratos de gas

Preguntas científicas

- Cinética de la formación y descomposición de clatratos
- Estudio de las propiedades físico-químicas de los clatratos bajo condiciones de las cortezas de los satélites de hielo
- Rango de estabilidad de clatratos, incluyendo las posibles transiciones de fase que cambiarían sus propiedades físicas y químicas

Metas específicas

- Síntesis de clatratos en celdas de gran volumen diseñadas específicamente
- Raman, IR y ultrasonidos
- Propiedades ópticas, eléctricas y dieléctricas
- Formación de clatratos desde salmueras a presiones del océano de Europa
- Determinación de diagramas PTx

I. AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

b. Efectos de alta presión sobre soluciones acuosas: sales, agregados supramoleculares y proteínas

Preguntas científicas:

- Efectos de la presión en la constante de equilibrio de las soluciones acuosas
- Efectos de la presión en las propiedades físico-químicas de los hidratos en las condiciones de las cortezas de los satélites de hielo
- Experimentos y simulaciones de soluciones de proteínas

Metas específicas:

- Sistemas acuosos con sulfatos de Mg^{2+} , Na^{+} , $Fe^{2+}/3^{+}$; los sistemas multi-componente son de interés para las ciencias de la Tierra y del Espacio.
- Diagramas pH, eH en función de la presión
- Propiedades reológicas y termodinámicas (densidad y coeficientes de expansión) de fases ultra-hidratadas of ultra hydrated phases of these ions will be studied to several GPa.
- Pressure effects on cyclodextrine-based host-guest systems
- Micelle formation and stability: caseine and other systems of interest in food science and technology
- Studying the conformational changes in proteins under pressure

I. AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

c. Microbiología en condiciones extremas de presión y temperatura

Preguntas científicas

- Adaptación de los organismos vivos bajo presión como campo de interés de biología y tecnología de los alimentos
- Identificación de los factores clave que determinan la piezorresistencia en arqueas y eubacterias extremófilas, especialmente la piezorresistencia de proteínas y los genes relacionados
- Influencia de los factores previos en el origen de la vida (básica) y procesamiento de alimentos (aplicado)

Metas específicas

- Estudio de procesos de oxidación lipídica en membranas celulares bajo presión
- Estudio de integridad y funcionalidad de las membranas celulares bajo presión
- Modelización de comportamiento de las membranas celulares bajo presión como liposomas de composición simple
- Identificación de proteínas responsables de piezoresistencia y genes relacionados

Clatratos en Europa


Master de tesis de Vicky – Octubre 2010
Clatratos en Europa, satélite de Júpiter

En el sistema solar exterior se dan las condiciones y los materiales necesarios para que se formen clatratos de gas, tanto en los satélites de hielo como en los cometas

- agua
- volátiles
- P y T

Manifestaciones geológicas observadas en los satélites de hielo pueden ser producto de la descomposición de los clatratos de gas

- estructuras de terrenos caóticos
- actividad de géiseres



EUROPA

Europa es un satélite especialmente interesante para Astrobiología debido a que parece albergar en su interior un océano de agua líquida en contacto con un substrato rocoso.

Actualmente Europa tiene una superficie de hielo dominada por H_2O , sales hidratadas y/o hidratos de ácido sulfúrico, CO_2 , SO_2 y H_2S .

El océano de agua líquida está en contacto con la capa rocosa probablemente afectada por actividad hidrotermal, cuyas fumarolas liberan gases.

Clatratos de gases volcánicos en el interior. De oxidantes radiogénicos en la subsuperficie.

APLICACIONES:

- Terrenos caóticos
- Diferenciación criomagmática



www.malta-consolider.com

M1 CONSOLIDER
ALTA

Matter at High Pressure

PREMIO
NACIONAL
2010

ESTABILIDAD CLATRATOS



CO₂ → Estabilidad desde 0.05-0.5 mm de profundidad hasta el océano




H₂S → Estabilidad desde 0.25-1.5 mm de profundidad hasta el océano





CLATRATOS VS. SALMUERAS

Tradicionalmente los electrolitos en solución se han descrito como inhibidores de formación de clatratos debido a:

- La fuerte atracción del agua hacia los electrolitos.
- El efecto “*salting-out*”: descenso de la solubilidad del gas en la salmuera debido al enlace del agua con los iones.

Inhibidores típicos: metanol, cloruros. Muy estudiados debido a sus aplicaciones técnicas y su aplicación en el análisis de inclusiones fluidas.





www.malta-consolider.com

M1 CONSOLIDER
ALTA

Matter at High Pressure






2010

DIFERENCIACIÓN POR CLATRATIZACIÓN

La formación de clatratos desde un magma enriquecido en gases e iones disueltos puede producir la diferenciación de la cámara magmática en los satélites de hielo.

Durante el proceso, las sales disueltas pueden precipitar y separarse por su mayor densidad, permitiendo a la solución con gases cristalizar en clatratos de gas.

Si los clatratos se desestabilizan por descompresión, los criomagmas ricos en volátiles podrán ascender a la superficie, e incluso erupcionar explosivamente.



CO₂ en disoluciones acuosas planetarias



(desarrollo de Laura)

- CO₂ en soluciones sulfatadas a alta presión

Aplicación en criomagmatismo. Flotabilidad. Explosividad de las erupciones de criomagmas. Diferenciación por clatratación

- CO₂ y carbonatos

Aplicación en la formación de carbonatos en épocas tempranas de la Tierra y Marte



Carbonatos en Tierra y Marte primitivos

Los Carbonatos pueden ser precipitados inorgánicamente o ser biomedados

Las calizas en la Tierra desempeñan un papel muy importante en el ciclo del C, fijan el carbono y liberan CO₂

Actualmente los carbonatos precipitan en mares someros en donde el agua está cargada de CO₂ según la reacción:





Los carbonatos son geomarcadores de unas condiciones ambientales particulares. Y pueden ser biomarcadores

Existen periodos en la historia de la Tierra con carbonatos particulares ¿Cómo precipitan?


- Carbonatos arcaicos
- CAP Carbonates

¿Dónde están los carbonatos en Marte?

635 Ma cap-carbonate sequence, NW Namibia





Tierra primitiva-carbonatos vs. Alta P



Arcaico-atmósfera más densa. Hasta 200 bares de presión de CO₂

Proterozoico- los “cap carbonates” son calizas y dolomías que aparecen por encima de los depósitos glaciares proterozoicos (Makganyene ~ 2220 M.a., Sturtian ~700 M.a. o Marinoan 635 M.a.). Tienen características especiales en relación a carbonatos de otras épocas: cementos bien desarrollados, ondulaciones gigantes, barita diagenética, relación isotópica deficitaria en ¹³C etc.

Carbonatos Arcaicos



Los carbonatos de más de 2400 m.a. No son frecuentes en la Tierra. Son peculiares:



- Asociaciones de caliza y ortocuarcitas (sedimentarias o volcánicas)
- Carbonatos asociados a esquistos o cherts
- Carbonatos en asociaciones volcánicas
- Sedimentos clásticos-carbonatados

La atmósfera del Arcaico fue muy rica en CO_2 . Esto pudo acidificar los océanos

Los datos muestran que la alteración de las rocas ígneas liberó Ca^{2+} a los océanos

La corteza era delgada e inestable, con pocos lugares aptos para la precipitación masiva de carbonatos. Solo en partes profundas con favorable contenido en CO_2 (fáciles de destruir posteriormente por tectónica)

Carbonatos Proterozoicos. *Cap carbonates*



Carbonatos que están en contacto limpio con facies periglaciares. Son dolomias microcristalinas que aparecen por todo el globo, con calizas post-glaciares por encima

Se han propuesto 3 modelos que relacionan una deglaciación abrupta y el evento de precipitación de carbonatos (alcalinidad extrema)

- **cambio brusco de los océanos** anóxicos profundos, desgasificación de CO₂, subsecuente liberación de la alcalinidad en nuevas cuencas subsumergidas y precipitación de CaCO₃

- **Snow Ball Earth:** alto grado de alteración química debido a condiciones de un super efecto invernadero seguidas de una glaciación global. Deglaciación catastrófica cuando el nivel de pCO₂ fue crítico (0.12 bares)





- liberación masiva de la alcalinidad por desestabilización **de clatratos de metano**, debido a la oxidación anaeróbica por sulfato-reducción

Experimentos de precipitación de carbonatos para determinar las condiciones en las que ocurrió: P, T, acidez, redox??? Validación de alguna de las hipótesis mencionadas. **La dolomita no cristaliza directamente en columnas de agua en condiciones de la superficie terrestre.**

Marte primitivo-carbonatos vs. Alta P

Si Marte tuvo agua y una atmósfera mas densa podría haber formado carbonatos

El rover Spirit ha identificado carbonatos Mg-Fe (16 - 34 peso%) en Columbia Hills del cráter Gusev. Podrían haber precipitado de soluciones carbonatadas en condiciones hidrotermales a pH casi neutro en asociación con la actividad volcánica del periodo Noico



Modelling High Pressure Processes in Food Engineering

Nadia Smith, Ángel Manuel Ramos del Olmo

Research Group MOMAT - Universidad Complutense de Madrid - Spain
<http://www.mat.ucm.es/momat>



QUIMAPRES



Reunión MALTA-CONSOLIDER 2011

Madrid, January 24th, 2010

Consumers needs

Consumers want food in **good conditions**.

- **Problem:** Food **decomposition** due to microorganisms and enzymes.
- **Goal:** Prolong food **shelf life**, preserving **organoleptic** (flavour, colour, etc...) and **nutritional properties**.
- **Minimally processed** and **additive-free** products.

→ **Food processing**

- **Classical Industrial Methods:** Freezing, Sterilization, Pasteurization...
- Disadvantages: Loss of nutritional properties, change in flavour, oxidation (for process that use high temperatures) and non homogeneous crystallization (for classical freezing methods).

Consumers want food in **good conditions**.

- **Problem:** Food **decomposition** due to microorganisms and enzymes.
- **Goal:** Prolong food **shelf life**, preserving **organoleptic** (flavour, colour, etc...) and **nutritional properties**.
- **Minimally processed** and **additive-free** products.

→ **Food processing**

- **Classical Industrial Methods:** Freezing, Sterilization, Pasteurization...
- **Disadvantages:** Loss of nutritional properties, change in flavour, oxidation (for process that use high temperatures) and non homogeneous crystallization (for classical freezing methods).

Nowadays technology

New emerging technology → **High Pressure (HP) processes**

- Combination of HP and moderate temperatures inactivates microorganisms and enzymes preserving many nutritional and organoleptic properties.
- additive-free products!
- HP can also be used for freezing: uniform nucleation and crystallization

Our goal as mathematicians

Modelling of different kinds of HP processes for **simulation** and **optimization** purposes

- Mathematical modelling of enzymatic and microbial inactivation
- Heat Transfer modelling for HP-moderate temperature process and also for HP-freezing processes.

Nowadays technology

New emerging technology → **High Pressure (HP) processes**

- Combination of HP and moderate temperatures inactivates microorganisms and enzymes preserving many nutritional and organoleptic properties.
- additive-free products!
- HP can also be used for freezing: uniform nucleation and crystallization

Our goal as mathematicians

Modelling of different kinds of HP processes for **simulation** and **optimization** purposes

- Mathematical modelling of enzymatic and microbial inactivation
- Heat Transfer modelling for HP-moderate temperature process and also for HP-freezing processes.

Nowadays technology

New emerging technology → **High Pressure (HP) processes**

- Combination of HP and moderate temperatures inactivates microorganisms and enzymes preserving many nutritional and organoleptic properties.
- additive-free products!
- HP can also be used for freezing: uniform nucleation and crystallization

Our goal as mathematicians

Modelling of different kinds of HP processes for **simulation** and **optimization** purposes

- Mathematical modelling of enzymatic and microbial inactivation
- Heat Transfer modelling for HP-moderate temperature process and also for HP-freezing processes.

Nowadays technology

New emerging technology → **High Pressure (HP) processes**

- Combination of HP and moderate temperatures inactivates microorganisms and enzymes preserving many nutritional and organoleptic properties.
- additive-free products!
- HP can also be used for freezing: uniform nucleation and crystallization

Our goal as mathematicians

Modelling of different kinds of HP processes for **simulation** and **optimization** purposes

- Mathematical modelling of enzymatic and microbial inactivation
- Heat Transfer modelling for HP-moderate temperature process and also for HP-freezing processes.

Modelling HP-T on microorganisms and enzymes

Model for enzymatic inactivation

Definition (Inactivation)

Reduction of undesired biological activity, such as enzymatic catalysis and microbial contamination.

The enzymatic activity of a particular enzyme can usually be described by the first-order kinetic model

$$\begin{cases} \frac{dA(t; T, P)}{dt} = -k(T, P)A(t; T, P) \\ A(0; T, P) = A_0 \end{cases}$$

$A(t; T, P)$: enzymatic activity at time t , when the food sample follows a process at temperature T and pressure P ; A_0 : initial enzy. act.; k : *inactivation rate*

- Isothermal/Isobaric process $\Rightarrow A(t; T, P) = A_0 \exp(-k(T, P)t)$
- General process $\Rightarrow A(t; T, P) = A_0 \exp(-\int_0^t k(T(s), P(s))ds)$

Modelling HP-T on microorganisms and enzymes

Model for enzymatic inactivation

Definition (Inactivation)

Reduction of undesired biological activity, such as enzymatic catalysis and microbial contamination.

The enzymatic activity of a particular enzyme can usually be described by the first-order kinetic model

$$\begin{cases} \frac{dA(t; T, P)}{dt} = -k(T, P)A(t; T, P) \\ A(0; T, P) = A_0 \end{cases}$$

$A(t; T, P)$: enzymatic activity at time t , when the food sample follows a process at temperature T and pressure P ; A_0 : initial enzy. act.; k : *inactivation rate*

- Isothermal/Isobaric process $\Rightarrow A(t; T, P) = A_0 \exp(-k(T, P)t)$
- General process $\Rightarrow A(t; T, P) = A_0 \exp(-\int_0^t k(T(s), P(s))ds)$

Modelling HP-T on microorganisms and enzymes

Model for enzymatic inactivation

Definition (Inactivation)

Reduction of undesired biological activity, such as enzymatic catalysis and microbial contamination.

The enzymatic activity of a particular enzyme can usually be described by the first-order kinetic model

$$\begin{cases} \frac{dA(t; T, P)}{dt} = -k(T, P)A(t; T, P) \\ A(0; T, P) = A_0 \end{cases}$$

$A(t; T, P)$: enzymatic activity at time t , when the food sample follows a process at temperature T and pressure P ; A_0 : initial enzy. act.; k : *inactivation rate*

- Isothermal/Isobaric process $\Rightarrow A(t; T, P) = A_0 \exp(-k(T, P)t)$
- General process $\Rightarrow A(t; T, P) = A_0 \exp(-\int_0^t k(T(s), P(s))ds)$

Modelling HP-T on microorganisms and enzymes

Model for bacterial inactivation

To describe the evolution of a microbial population, we can follow the above model, just by changing enzymatic activity $A(t; T, P)$ by microbial population $N(t; T, P)$.

Or this other model

$$\begin{cases} \frac{dN(t; T, P)}{dt} = -\frac{\ln(10)}{D(T, P)} N(t; T, P) \\ N(0; T, P) = N_0 \end{cases}$$

$D(T, P)$: decimal reduction time, or time required for a 1-log-cycle reduction in the microbial population.

- Isothermal/Isobaric process $\Rightarrow N(t; T, P) = N_0 10^{-\frac{t}{D(T, P)}}$
- General process $\Rightarrow N(t; T, P) = N_0 10^{-\int_0^t \frac{1}{D(T(s), P(s))} ds}$

Modelling HP-T on microorganisms and enzymes

Model for bacterial inactivation

To describe the evolution of a microbial population, we can follow the above model, just by changing enzymatic activity $A(t; T, P)$ by microbial population $N(t; T, P)$.

Or this other model

$$\begin{cases} \frac{dN(t; T, P)}{dt} = -\frac{\ln(10)}{D(T, P)} N(t; T, P) \\ N(0; T, P) = N_0 \end{cases}$$

$D(T, P)$: *decimal reduction time*, or time required for a 1-log-cycle reduction in the microbial population.

- Isothermal/Isobaric process $\Rightarrow N(t; T, P) = N_0 10^{-\frac{t}{D(T, P)}}$
- General process $\Rightarrow N(t; T, P) = N_0 10^{-\int_0^t \frac{1}{D(T(s), P(s))} ds}$

Modelling HP-T on microorganisms and enzymes

Model for bacterial inactivation

To describe the evolution of a microbial population, we can follow the above model, just by changing enzymatic activity $A(t; T, P)$ by microbial population $N(t; T, P)$.

Or this other model

$$\begin{cases} \frac{dN(t; T, P)}{dt} = -\frac{\ln(10)}{D(T, P)} N(t; T, P) \\ N(0; T, P) = N_0 \end{cases}$$

$D(T, P)$: *decimal reduction time*, or time required for a 1-log-cycle reduction in the microbial population.

- Isothermal/Isobaric process $\Rightarrow N(t; T, P) = N_0 10^{-\frac{t}{D(T, P)}}$
- General process $\Rightarrow N(t; T, P) = N_0 10^{-\int_0^t \frac{1}{D(T(s), P(s))} ds}$

Modelling HP-T on microorganisms and enzymes

Model for bacterial inactivation

To describe the evolution of a microbial population, we can follow the above model, just by changing enzymatic activity $A(t; T, P)$ by microbial population $N(t; T, P)$.

Or this other model

$$\begin{cases} \frac{dN(t; T, P)}{dt} = -\frac{\ln(10)}{D(T, P)} N(t; T, P) \\ N(0; T, P) = N_0 \end{cases}$$

$D(T, P)$: *decimal reduction time*, or time required for a 1-log-cycle reduction in the microbial population.

- Isothermal/Isobaric process $\Rightarrow N(t; T, P) = N_0 10^{-\frac{t}{D(T, P)}}$
- General process $\Rightarrow N(t; T, P) = N_0 10^{-\int_0^t \frac{1}{D(T(s), P(s))} ds}$

Identification of kinetic parameters

Identification of the inactivation rate constants $k(T, P)$ and $D(T, P)$

For general (T, P) -dependent processes we may use

- A combination of Eyring's and Arrhenius' equations:

$$k(T, P) = k_{T_{\text{ref}}, P_{\text{ref}}} \exp \left(-B \left(\frac{1}{T} - \frac{1}{T_{\text{ref}}} \right) \right) \exp(-C(P - P_{\text{ref}}))$$

- A more complex choice given by

$$k(P, T) = k_{\text{ref}} \exp \left(\frac{-\Delta V_{\text{ref}}}{RT} (P - P_{\text{ref}}) \right) \exp \left(\frac{\Delta S_{\text{ref}}}{RT} (T - T_{\text{ref}}) \right) \\ \exp \left(\frac{-\Delta \kappa}{2RT} (P - P_{\text{ref}})^2 \right) \exp \left(\frac{-2\Delta \zeta}{RT} (P - P_{\text{ref}})(T - T_{\text{ref}}) \right) \\ \exp \left(\frac{\Delta C_p}{RT} \left(T \left(\ln \frac{T}{T_{\text{ref}}} - 1 \right) + T_{\text{ref}} \right) \right) + \text{high order terms,}$$

- For the microbial decimal reduction time $D(T, P)$ we have not found any formula in the literature. By construction $k = \frac{\ln(10)}{D}$

Identification of kinetic parameters

Identification of the inactivation rate constants $k(T, P)$ and $D(T, P)$

For general (T, P) –dependent processes we may use

- A combination of Eyring's and Arrhenius' equations:

$$k(T, P) = k_{T_{\text{ref}}, P_{\text{ref}}} \exp\left(-B\left(\frac{1}{T} - \frac{1}{T_{\text{ref}}}\right)\right) \exp(-C(P - P_{\text{ref}}))$$

- A more complex choice given by

$$k(P, T) = k_{\text{ref}} \exp\left(\frac{-\Delta V_{\text{ref}}}{RT}(P - P_{\text{ref}})\right) \exp\left(\frac{\Delta S_{\text{ref}}}{RT}(T - T_{\text{ref}})\right) \\ \exp\left(\frac{-\Delta \kappa}{2RT}(P - P_{\text{ref}})^2\right) \exp\left(\frac{-2\Delta \zeta}{RT}(P - P_{\text{ref}})(T - T_{\text{ref}})\right) \\ \exp\left(\frac{\Delta C_p}{RT}\left(T\left(\ln\frac{T}{T_{\text{ref}}} - 1\right) + T_{\text{ref}}\right)\right) + \text{high order terms,}$$

- For the microbial decimal reduction time $D(T, P)$ we have not found any formula in the literature. By construction $k = \frac{\ln(10)}{D}$

Identification of kinetic parameters

Identification of the inactivation rate constants $k(T, P)$ and $D(T, P)$

For general (T, P) –dependent processes we may use

- A combination of Eyring's and Arrhenius' equations:

$$k(T, P) = k_{T_{\text{ref}}, P_{\text{ref}}} \exp\left(-B\left(\frac{1}{T} - \frac{1}{T_{\text{ref}}}\right)\right) \exp(-C(P - P_{\text{ref}}))$$

- A more complex choice given by

$$k(P, T) = k_{\text{ref}} \exp\left(\frac{-\Delta V_{\text{ref}}}{RT}(P - P_{\text{ref}})\right) \exp\left(\frac{\Delta S_{\text{ref}}}{RT}(T - T_{\text{ref}})\right) \\ \exp\left(\frac{-\Delta \kappa}{2RT}(P - P_{\text{ref}})^2\right) \exp\left(\frac{-2\Delta \zeta}{RT}(P - P_{\text{ref}})(T - T_{\text{ref}})\right) \\ \exp\left(\frac{\Delta C_p}{RT}\left(T\left(\ln\frac{T}{T_{\text{ref}}}\right) - 1\right) + T_{\text{ref}}\right) + \text{high order terms,}$$

- For the microbial decimal reduction time $D(T, P)$ we have not found any formula in the literature. By construction $k = \frac{\ln(10)}{D}$

Solving the inactivation models

- We identify the parameters ($k_{T,P}$, T_{ref} , P_{ref} , $k_{T_{\text{ref}},P_{\text{ref}}}$, B , C , z_T , z_P , $\dots$) by using **experimental data** and **regression techniques**.
- We can now model enzymatic inactivation or microbial population for $HP-T$ processes, using the models for $A(t; T, P)$ or $N(t; T, P)$. For this you need **temperature** and **pressure** curves.
- The pressure is known a priori, but the temperature changes during the process (adiabatic heating effect, cooling or heating on the walls of the chamber, etc...)

→ **Heat Transfer models**

Solving the inactivation models

- We identify the parameters ($k_{T,P}$, T_{ref} , P_{ref} , $k_{T_{\text{ref}},P_{\text{ref}}}$, B , C , z_T , z_P , $\dots$) by using **experimental data** and **regression techniques**.
- We can now model enzymatic inactivation or microbial population for $HP-T$ processes, using the models for $A(t; T, P)$ or $N(t; T, P)$. For this you need **temperature** and **pressure** curves.
- The pressure is known a priori, but the temperature changes during the process (adiabatic heating effect, cooling or heating on the walls of the chamber, etc...)

→ **Heat Transfer models**

Solving the inactivation models

- We identify the parameters ($k_{T,P}$, T_{ref} , P_{ref} , $k_{T_{\text{ref}},P_{\text{ref}}}$, B , C , z_T , z_P , $\dots$) by using **experimental data** and **regression techniques**.
- We can now model enzymatic inactivation or microbial population for $HP-T$ processes, using the models for $A(t; T, P)$ or $N(t; T, P)$. For this you need **temperature** and **pressure** curves.
- The pressure is known a priori, but the temperature changes during the process (adiabatic heating effect, cooling or heating on the walls of the chamber, etc...)


→ **Heat Transfer models**

Solving the inactivation models


- We identify the parameters ($k_{T,P}$, T_{ref} , P_{ref} , $k_{T_{\text{ref}},P_{\text{ref}}}$, B , C , z_T , z_P , $\dots$) by using **experimental data** and **regression techniques**.
- We can now model enzymatic inactivation or microbial population for $HP-T$ processes, using the models for $A(t; T, P)$ or $N(t; T, P)$. For this you need **temperature** and **pressure** curves.
- The pressure is known a priori, but the temperature changes during the process (adiabatic heating effect, cooling or heating on the walls of the chamber, etc...)

→ **Heat Transfer models**

Graphical description of the HP Equipment




Graphical description of the HP Equipment



Graphical description of the HP Equipment

Computational domain

2D model (r,z) - Computational domain (big and small food sample)



Different heat transfer models

During a $HP-T$ process the temperature of the processed food may change with time and with space. There are many models that describe the evolution of temperature of the processed food

- **ODE models:** when temperature only depends on time

$$\frac{dT}{dt} = H(T^e - T) + \alpha T \frac{dP}{dt}$$

- **PDE models:** when temperature depends on time and space.
Different models depending on type of food sample

- **Heat transfer by conduction: For solid type foods (big sample)**
- Heat transfer by conduction and convection (only in the pressurizing fluid): For solid type foods (small sample)
- Heat transfer by conduction and convection (in food and fluid): For liquid type foods

Different heat transfer models

During a $HP-T$ process the temperature of the processed food may change with time and with space. There are many models that describe the evolution of temperature of the processed food

- **ODE models:** when temperature only depends on time

$$\frac{dT}{dt} = H(T^e - T) + \alpha T \frac{dP}{dt}$$

- **PDE models:** when temperature depends on time and space.
Different models depending on type of food sample

- **Heat transfer by conduction: For solid type foods (big sample)**
- Heat transfer by conduction and convection (only in the pressurizing fluid): For solid type foods (small sample)
- Heat transfer by conduction and convection (in food and fluid): For liquid type foods

Different heat transfer models

During a $HP-T$ process the temperature of the processed food may change with time and with space. There are many models that describe the evolution of temperature of the processed food

- **ODE models:** when temperature only depends on time

$$\frac{dT}{dt} = H(T^e - T) + \alpha T \frac{dP}{dt}$$

- **PDE models:** when temperature depends on time and space.
Different models depending on type of food sample


- **Heat transfer by conduction: For solid type foods (big sample)**
- Heat transfer by conduction and convection (only in the pressurizing fluid): For solid type foods (small sample)
- Heat transfer by conduction and convection (in food and fluid): For liquid type foods

Heat transfer by conduction

Solid type foods. Big sample

By using cylindrical coordinates and taking into account axial symmetry, our starting point is the **(r,z) 2D-model**

$$\left\{ \begin{array}{ll} \rho C_p \frac{\partial T}{\partial t} - \frac{1}{r} \frac{\partial}{\partial r} \left(r k \frac{\partial T}{\partial r} \right) - \frac{\partial}{\partial z} \left(k \frac{\partial T}{\partial z} \right) = \alpha \frac{dP}{dt} T & \text{on } (\Gamma \setminus (\Gamma_r \cup \Gamma_{up})) , \\ k \frac{\partial T}{\partial \mathbf{n}} = 0 & \\ k \frac{\partial T}{\partial \mathbf{n}} = h(T_{\text{env}} - T) & \text{on } \Gamma_{up}, \\ T = T_r & \text{on } \Gamma_r, \\ T(0) = T_0 & \text{in } \Omega. \end{array} \right.$$




→ Can we simplify this model?

Heat transfer by conduction

Solid type foods. Big sample

By using cylindrical coordinates and taking into account axial symmetry, our starting point is the **(r,z) 2D-model**

$$\left\{ \begin{array}{ll} \rho C_p \frac{\partial T}{\partial t} - \frac{1}{r} \frac{\partial}{\partial r} \left(r k \frac{\partial T}{\partial r} \right) - \frac{\partial}{\partial z} \left(k \frac{\partial T}{\partial z} \right) = \alpha \frac{dP}{dt} T & \text{on } (\Gamma \setminus (\Gamma_r \cup \Gamma_{up})) , \\ k \frac{\partial T}{\partial \mathbf{n}} = 0 & \\ k \frac{\partial T}{\partial \mathbf{n}} = h(T_{\text{env}} - T) & \text{on } \Gamma_{up}, \\ T = T_r & \text{on } \Gamma_r, \\ T(0) = T_0 & \text{in } \Omega. \end{array} \right.$$



→ **Can we simplify this model?**

Dimensional analysis

Non-dimensionalization (^ denote non-dimensional variables):

$$\hat{r} = \frac{r}{R}, \quad \hat{z} = \frac{z}{Z}, \quad \hat{t} = \frac{t}{\tau}, \quad \hat{T} = \frac{T - T_r}{\Delta T}$$

Consider

$$\frac{dP}{dt}(t) = \begin{cases} \frac{\gamma}{t_p}, & 0 < t \leq t_p, \\ 0, & t > t_p \end{cases}$$

Dropping the ^ symbol:

$$\frac{\partial T}{\partial t} - \frac{k\tau}{R^2 \rho C_p} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) - \frac{k\tau}{Z^2 \rho C_p} \frac{\partial^2 T}{\partial z^2} = \frac{\alpha \gamma \tau}{\rho C_p t_p} T + \frac{\alpha \gamma \tau}{\rho C_p t_p} \frac{T_r}{\Delta T}$$

Choose scales: R = width food sample, Z = height food sample,

$$\Delta T = \max\{|T_0 - T_r|, \frac{\alpha \gamma T_0}{\rho C_p}\}, \quad \tau = \frac{\rho C_p t_p}{\alpha \gamma} \min\left\{1, \frac{\Delta T}{T_r}\right\}$$

Dimensional analysis

Non-dimensionalization (^ denote non-dimensional variables):

$$\hat{r} = \frac{r}{R}, \quad \hat{z} = \frac{z}{Z}, \quad \hat{t} = \frac{t}{\tau}, \quad \hat{T} = \frac{T - T_r}{\Delta T}$$

Consider

$$\frac{dP}{dt}(t) = \begin{cases} \frac{\gamma}{t_p}, & 0 < t \leq t_p, \\ 0, & t > t_p \end{cases}$$

Dropping the ^ symbol:

$$\frac{\partial T}{\partial t} - \frac{k\tau}{R^2 \rho C_p} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) - \frac{k\tau}{Z^2 \rho C_p} \frac{\partial^2 T}{\partial z^2} = \frac{\alpha \gamma \tau}{\rho C_p t_p} T + \frac{\alpha \gamma \tau}{\rho C_p t_p} \frac{T_r}{\Delta T}$$

Choose scales: R = width food sample, Z = height food sample,

$$\Delta T = \max\{|T_0 - T_r|, \frac{\alpha \gamma T_0}{\rho C_p}\}, \quad \tau = \frac{\rho C_p t_p}{\alpha \gamma} \min\left\{1, \frac{\Delta T}{T_r}\right\}$$

Dimensional analysis

Non-dimensionalization (^ denote non-dimensional variables):

$$\hat{r} = \frac{r}{R}, \quad \hat{z} = \frac{z}{Z}, \quad \hat{t} = \frac{t}{\tau}, \quad \hat{T} = \frac{T - T_r}{\Delta T}$$

Consider

$$\frac{dP}{dt}(t) = \begin{cases} \frac{\gamma}{t_p}, & 0 < t \leq t_p, \\ 0, & t > t_p \end{cases}$$

Dropping the ^ symbol:

$$\frac{\partial T}{\partial t} - \frac{k\tau}{R^2 \rho C_p} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) - \frac{k\tau}{Z^2 \rho C_p} \frac{\partial^2 T}{\partial z^2} = \frac{\alpha \gamma \tau}{\rho C_p t_p} T + \frac{\alpha \gamma \tau}{\rho C_p t_p} \frac{T_r}{\Delta T}$$

Choose scales: R = width food sample, Z = height food sample,

$$\Delta T = \max\{|T_0 - T_r|, \frac{\alpha \gamma T_0}{\rho C_p}\}, \quad \tau = \frac{\rho C_p t_p}{\alpha \gamma} \min\left\{1, \frac{\Delta T}{T_r}\right\}$$

Dimensional analysis

Non-dimensionalization (^ denote non-dimensional variables):

$$\hat{r} = \frac{r}{R}, \quad \hat{z} = \frac{z}{Z}, \quad \hat{t} = \frac{t}{\tau}, \quad \hat{T} = \frac{T - T_r}{\Delta T}$$

Consider

$$\frac{dP}{dt}(t) = \begin{cases} \frac{\gamma}{t_p}, & 0 < t \leq t_p, \\ 0, & t > t_p \end{cases}$$

Dropping the ^ symbol:

$$\frac{\partial T}{\partial t} - \frac{k\tau}{R^2 \rho C_p} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) - \frac{k\tau}{Z^2 \rho C_p} \frac{\partial^2 T}{\partial z^2} = \frac{\alpha \gamma \tau}{\rho C_p t_p} T + \frac{\alpha \gamma \tau}{\rho C_p t_p} \frac{T_r}{\Delta T}$$

Choose scales: R = width food sample, Z = height food sample,

$$\Delta T = \max\{|T_0 - T_r|, \frac{\alpha \gamma T_0}{\rho C_p}\}, \quad \tau = \frac{\rho C_p t_p}{\alpha \gamma} \min\left\{1, \frac{\Delta T}{T_r}\right\}$$

Dimensional analysis

Non-dimensionalization (^ denote non-dimensional variables):

$$\hat{r} = \frac{r}{R}, \quad \hat{z} = \frac{z}{Z}, \quad \hat{t} = \frac{t}{\tau}, \quad \hat{T} = \frac{T - T_r}{\Delta T}$$

Consider

$$\frac{dP}{dt}(t) = \begin{cases} \frac{\gamma}{t_p}, & 0 < t \leq t_p, \\ 0, & t > t_p \end{cases}$$

Dropping the ^ symbol:

$$\frac{\partial T}{\partial t} - \underbrace{\frac{k\tau}{R^2 \rho C_p}}_a \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) - \underbrace{\frac{k\tau}{Z^2 \rho C_p}}_b \frac{\partial^2 T}{\partial z^2} = \underbrace{\frac{\alpha \gamma \tau}{\rho C_p t_p}}_c T + \underbrace{\frac{\alpha \gamma \tau}{\rho C_p t_p} \frac{T_r}{\Delta T}}_d$$

Choose scales: R = width food sample, Z = height food sample,

$$\Delta T = \max\{|T_0 - T_r|, \frac{\alpha \gamma T_0}{\rho C_p}\}, \quad \tau = \frac{\rho C_p t_p}{\alpha \gamma} \min\left\{1, \frac{\Delta T}{T_r}\right\}$$

Non-dimensional system

With all the previously chosen scales, we can now rewrite the system in a simplified non-dimensional form:

$$\left\{ \begin{array}{ll} \frac{\partial T}{\partial t} - \mathbf{a} \frac{1}{r} \frac{\partial}{\partial r} \left(\frac{\partial T}{\partial r} \right) - \mathbf{b} \frac{\partial^2 T}{\partial z^2} = (\mathbf{c}T + \mathbf{d})\chi_{(0, \frac{t_p}{\tau})}(t) & \text{in } \hat{\Omega} \\ \frac{\partial T}{\partial \mathbf{n}} = 0 & \text{on } (\hat{\Gamma} \setminus (\hat{\Gamma}_r \cup \hat{\Gamma}_{\text{up}})) \\ \frac{\partial T}{\partial \mathbf{n}} = -\frac{hZ}{k} T & \text{on } \hat{\Gamma}_{\text{up}}, \\ T = 0 & \text{on } \hat{\Gamma}_r, \\ T(0) = \frac{T_0 - T_r}{\Delta T} & \text{in } \hat{\Omega}. \end{array} \right.$$

The values of the parameters **a**, **b**, **c**, **d** depend on the food sample, the pressurizing fluid, the dimensions of the holder, the pressure profile, the process time, etc...

Example of a HP process

The pressure profile is given by **Process P** (360 MPa):

$$T_0 = 40^\circ\text{C} \text{ in } \Omega, \quad \frac{dP}{dt} = \begin{cases} \frac{120}{61} \text{ MPa s}^{-1}, & 0 < t \leq 183 \\ 0 \text{ MPa s}^{-1}, & t > 183. \end{cases}$$

Using thermophysical data of **tylose** as the food simile, **water** as the pressurizing medium, and for a **900 seconds** process, the values of the parameters are

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
$\hat{\Omega}_F$	0.02	0.00126	0.07	1
$\hat{\Omega}_P$	0.023	0.0014	0.05	0.7
$\hat{\Omega}_C$	0.013	0.00081	0	0
$\hat{\Omega}_S$	2.423	0.148	0	0

Key point: Conductivity height coefficient (**b**) was smaller than the radial one.

Is it reasonable to neglect that term? $\longrightarrow$ **1D model?**

Example of a HP process

The pressure profile is given by **Process P** (360 MPa):

$$T_0 = 40^{\circ}\text{C} \text{ in } \Omega, \quad \frac{dP}{dt} = \begin{cases} \frac{120}{61} \text{ MPa s}^{-1}, & 0 < t \leq 183 \\ 0 \text{ MPa s}^{-1}, & t > 183. \end{cases}$$

Using thermophysical data of **tylose** as the food simile, **water** as the pressurizing medium, and for a **900 seconds** process, the values of the parameters are

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
$\hat{\Omega}_F$	0.02	0.00126	0.07	1
$\hat{\Omega}_P$	0.023	0.0014	0.05	0.7
$\hat{\Omega}_C$	0.013	0.00081	0	0
$\hat{\Omega}_S$	2.423	0.148	0	0

Key point: Conductivity height coefficient (**b**) was smaller than the radial one.

Is it reasonable to neglect that term? $\longrightarrow$ **1D model?**

Example of a HP process

The pressure profile is given by **Process P** (360 MPa):

$$T_0 = 40^\circ\text{C} \text{ in } \Omega, \quad \frac{dP}{dt} = \begin{cases} \frac{120}{61} \text{ MPa s}^{-1}, & 0 < t \leq 183 \\ 0 \text{ MPa s}^{-1}, & t > 183. \end{cases}$$

Using thermophysical data of **tylose** as the food simile, **water** as the pressurizing medium, and for a **900 seconds** process, the values of the parameters are

	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
$\hat{\Omega}_F$	0.02	0.00126	0.07	1
$\hat{\Omega}_P$	0.023	0.0014	0.05	0.7
$\hat{\Omega}_C$	0.013	0.00081	0	0
$\hat{\Omega}_S$	2.423	0.148	0	0

Key point: Conductivity height coefficient (**b**) was smaller than the radial one.

Is it reasonable to neglect that term? $\longrightarrow$ 1D model?

Example of a HP process

The pressure profile is given by **Process P** (360 MPa):

$$T_0 = 40^\circ\text{C} \text{ in } \Omega, \quad \frac{dP}{dt} = \begin{cases} \frac{120}{61} \text{ MPa s}^{-1}, & 0 < t \leq 183 \\ 0 \text{ MPa s}^{-1}, & t > 183. \end{cases}$$

Using thermophysical data of **tylose** as the food simile, **water** as the pressurizing medium, and for a **900 seconds** process, the values of the parameters are



	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>
$\hat{\Omega}_F$	0.02	0.00126	0.07	1
$\hat{\Omega}_P$	0.023	0.0014	0.05	0.7
$\hat{\Omega}_C$	0.013	0.00081	0	0
$\hat{\Omega}_S$	2.423	0.148	0	0

Key point: Conductivity height coefficient (**b**) was smaller than the radial one.

Is it reasonable to neglect that term? $\longrightarrow$ **1D model?**

Graphical results of the 2D-model

Solution of 2D model inside the food during process P at different heights and different radius



1D Heat transfer model

$$\left\{ \begin{array}{ll} \frac{\partial T}{\partial t} - a \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) = (c T + d) \chi_{(0,0,56)}(t) & \text{in } (0, 2) \times (0, 2,76), \\ \frac{\partial T}{\partial r} = 0 & \text{on } \{0\} \times (0, 2,76), \\ T = 0 & \text{on } \{2\} \times (0, 2,76), \\ T(0) = 1 & \text{in } (0, 2), \end{array} \right.$$

- This model is much faster to solve computationally than the 2D-model.
- But even better: we can find an **analytic solution** for the temperature inside the food!

$$T_F(r, t) = \begin{cases} t + 1, & t \leq 0,56, \\ \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) \exp(-0,02 \lambda_n^2 (t - 0,56)), & t > 0,56, \end{cases}$$

1D Heat transfer model

$$\left\{ \begin{array}{ll} \frac{\partial T}{\partial t} - a \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) = (c T + d) \chi_{(0,0,56)}(t) & \text{in } (0, 2) \times (0, 2,76), \\ \frac{\partial T}{\partial r} = 0 & \text{on } \{0\} \times (0, 2,76), \\ T = 0 & \text{on } \{2\} \times (0, 2,76), \\ T(0) = 1 & \text{in } (0, 2), \end{array} \right.$$

- This model is much faster to solve computationally than the 2D-model.
- But even better: we can find an **analytic solution** for the temperature inside the food!

$$T_F(r, t) = \begin{cases} t + 1, & t \leq 0,56, \\ \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) \exp(-0,02 \lambda_n^2 (t - 0,56)), & t > 0,56, \end{cases}$$

1D Heat transfer model

$$\left\{ \begin{array}{ll} \frac{\partial T}{\partial t} - a \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) = (c T + d) \chi_{(0,0,56)}(t) & \text{in } (0, 2) \times (0, 2,76), \\ \frac{\partial T}{\partial r} = 0 & \text{on } \{0\} \times (0, 2,76), \\ T = 0 & \text{on } \{2\} \times (0, 2,76), \\ T(0) = 1 & \text{in } (0, 2), \end{array} \right.$$

- This model is much faster to solve computationally than the 2D-model.
- But even better: we can find an **analytic solution** for the temperature inside the food!

$$T_F(r, t) = \begin{cases} t + 1, & t \leq 0,56, \\ \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) \exp(-0,02 \lambda_n^2 (t - 0,56)), & t > 0,56, \end{cases}$$


1D Heat transfer model

$$\left\{ \begin{array}{ll} \frac{\partial T}{\partial t} - a \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial T}{\partial r} \right) = (c T + d) \chi_{(0,0,56)}(t) & \text{in } (0, 2) \times (0, 2,76), \\ \frac{\partial T}{\partial r} = 0 & \text{on } \{0\} \times (0, 2,76), \\ T = 0 & \text{on } \{2\} \times (0, 2,76), \\ T(0) = 1 & \text{in } (0, 2), \end{array} \right.$$

- This model is much faster to solve computationally than the 2D-model.
- But even better: we can find an **analytic solution** for the temperature inside the food!

$$T_F(r, t) = \begin{cases} t + 1, & t \leq 0,56, \\ \sum_{n=1}^{\infty} C_n J_0(\lambda_n r) \exp(-0,02 \lambda_n^2 (t - 0,56)), & t > 0,56, \end{cases}$$


Pros and cons of the simplifications



Temperature differences between 2D and 1D model at the same radial points inside the food. **Mid height**. Process **P**.

We have tried with different pressure curves and conditions →
Very small differences for nearly every point inside food sample, **except**
for points very close to the top (**Rubber cap**) or bottom (**Steel**) →
Studying possibility of **boundary layers**. Working on **asymptotic expansion** right now.


Pros and cons of the simplifications



Temperature differences between 2D and 1D model at the same radial points inside the food. **Mid height**. Process **P**.

We have tried with different pressure curves and conditions →
Very small differences for nearly every point inside food sample, **except**
for points very close to the top (**Rubber cap**) or bottom (**Steel**) →
Studying possibility of **boundary layers**. Working on **asymptotic expansion** right now.

Pros and cons of the simplifications



Temperature differences between 2D and 1D model at the same radial points inside the food. **Mid height**. Process **P**.

We have tried with different pressure curves and conditions →
Very small differences for nearly every point inside food sample, **except**
for points very close to the top (**Rubber cap**) or bottom (**Steel**) →
Studying possibility of **boundary layers**. Working on **asymptotic expansion** right now.

Coupling of Enzymatic Inactivation and Heat Transfer Simplified Models



We can now couple the heat transfer models with kinetic equation for enzymatic activity $A(t; T, P)$. We compare the **resulting enzymatic activity** distribution that we obtain using the 2D heat transfer models versus the 1D models.

Model (for T)	BSAA	LOX	CPE
1D	26.92	68.66	93.88
2D-mid height	25.77	67.06	93.76
2D-full	26.43	67.18	93.86

Mean activity at final time for the 3 models considered and for BSAA (*Bacillus Subtilis* α -Amylase), LOX (Lipoxygenase) and CPE (Carrot Pectin Methyl-Esterase) enzymes.

- J. A. Infante, B. Ivorra, Á. M. Ramos and J. M. Rey, *On the Modelling and Simulation of High Pressure Processes and Inactivation of Enzymes in Food Engineering*. Mathematical Models and Methods in Applied Sciences (M3AS), Vol. 19, No. 12, (2009) pp. 2203–2229.

Modelling freezing processes with High pressure. HPSF processes



We are developing a heat transfer model for HPSF processes with **rapid** (a) or **slow** (b) expansions. For rapid expansion the **phase transition** occurs at **atmospheric pressure** conditions and for slow expansions **under pressure**.

The enthalpy formulation

Voller and Swaminathan (1990): g_l (liquid volume fraction) and g_s (solid volume fraction). "Mixture enthalpy":

$$e = g_s \int_{T_{\text{ref}}}^T \rho_s C_{p_s} d\theta + g_l \int_{T_{\text{ref}}}^T \rho_l C_{p_l} d\theta + \rho_l g_l \lambda,$$

where T_{ref} is a reference temperature (it is the temperature at which all the latent heat has been released) and λ is the latent heat.

"Mixture conductivity":

$$k = g_s k_s + g_l k_l.$$

Energy equation in enthalpy formulation

$$\frac{\partial e}{\partial t} = \nabla \cdot (k \nabla T).$$

Defining $C_{\text{vol}} = g_s \rho_s C_{p_s} + g_l \rho_l C_{p_l}$, and $\delta e = \int_{T_{\text{ref}}}^T (\rho_l C_{p_l} - \rho_s C_{p_s}) d\theta + \rho_l \lambda$,

$$\frac{\partial e}{\partial t} = C_{\text{vol}} \frac{\partial T}{\partial t} + \delta e \frac{\partial g_l}{\partial t}.$$

The enthalpy formulation: Source based method

Now it is possible to give the equation in a formulation for temperature:

$$C_{\text{vol}} \frac{\partial T}{\partial t} - \nabla \cdot (k \nabla T) = \alpha \frac{dP}{dt} T - \delta e \frac{\partial g_l}{\partial t}$$

More convenient: only one variable and easier to adapt the algorithms to existing numerical codes.

Problems: How to calculate the liquid fraction as a function of time $g_l(t)$. How to release the exact amount of latent heat λ for the temperature to increase up to the corresponding freezing point at atmospheric conditions.

We assume that the phase transition occurs at atmospheric pressure conditions (rapid expansion) and over a short period of time.

The enthalpy formulation: Source based method

Now it is possible to give the equation in a formulation for temperature:

$$C_{\text{vol}} \frac{\partial T}{\partial t} - \nabla \cdot (k \nabla T) = \alpha \frac{dP}{dt} T - \delta e \frac{\partial g_l}{\partial t}$$

More convenient: only one variable and easier to adapt the algorithms to existing numerical codes.

Problems: How to calculate the liquid fraction as a function of time $g_l(t)$. How to release the exact amount of latent heat λ for the temperature to increase up to the corresponding freezing point at atmospheric conditions.

We assume that the phase transition occurs at atmospheric pressure conditions (rapid expansion) and over a short period of time.

The enthalpy formulation: Source based method

Now it is possible to give the equation in a formulation for temperature:

$$C_{\text{vol}} \frac{\partial T}{\partial t} - \nabla \cdot (k \nabla T) = \alpha \frac{dP}{dt} T - \delta e \frac{\partial g_l}{\partial t}$$


More convenient: only one variable and easier to adapt the algorithms to existing numerical codes.

Problems: How to calculate the liquid fraction as a function of time $g_l(t)$. How to release the exact amount of latent heat λ for the temperature to increase up to the corresponding freezing point at atmospheric conditions.

We assume that the phase transition occurs at atmospheric pressure conditions (rapid expansion) and over a short period of time.

Experimental results of HPSF

Pressure and temperature experimental evolution in a gelatin sample during a High-pressure shift freezing process.




- P. P. Fernández, L. Otero, B. Guignon, P. D. Sanz, *High-pressure shift freezing versus high-pressure assisted freezing: Effects on the microstructure of a food model*. Food Hydrocolloids 20 (2006), pp. 510–522.


Numerical results when modelling HPSF

Example of HPSF process with rapid expansion of agar gel.

Temperature Agar of 2d model for different heights (at center radius). HPSF



Temperature Agar of 2d model for different radius (at mid height). HPSF




Evolution of the temperature inside food during HPSF process for 2D model, for different heights (left) and radial points (right).


Numerical examples when modelling HPSF

We again perform dimensional analysis and try to go to a 1D model.

Different radius of 1d model compared to 2d model (at mid height). HPSF



Different heights of 2d model compared to 1d model (at central radius). HPSF



Difference of temperatures of 2D model and 1D model inside food during HPSF process for different heights (left) and radial points (right).

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- ... a lot of research is still to be done in this area

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- ... a lot of research is still to be done in this area

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- ... a lot of research is still to be done in this area

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- ... a lot of research is still to be done in this area

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- ... a lot of research is still to be done in this area

Concluding Remarks

- The combination of high pressure with moderate temperatures inactivates microorganisms and enzymes without damaging the food significantly.
- We are able to predict the bacterial and enzymatic inactivation for high pressure processes in foods.
- These models are a very useful tool for equipment design and for optimization of the inactivation processes.
- Simplification of the models is convenient for quick simulations and for optimization purposes.
- Analytic solutions allow a qualitative study of the temperature profile
- **... a lot of research is still to be done in this area**

Emerging Risks in the Food Supply Chain

claud.j.cheftel@orange.fr

**Honorary Professor of Food Biochemistry & Technology
Université de Montpellier, France**

www.cheftel.com

**MALTA-CONSOLIDER
Miraflores de la Sierra, 24 Jan. 2011**

Emerging Risks

An **emerging risk** can be defined as (EFSA, 2007):

- a risk resulting from a **newly identified hazard** to which a significant exposure may occur
- or from an unexpected increased significant exposure and/or susceptibility **to a known hazard**.

Main Categories of Food Hazards & Risks

Types of Hazards

Manifestations

Populations at risk

**Pathogenic
Micro-organisms**

**Acute,
Immediate**

General
(individual & group
vulnerability)

Toxic chemicals

**Mainly chronic diseases
after prolonged exposure**
(often masked by influence
of other factors)

General
(individual & group
vulnerability)

Food allergens

Acute, Immediate
(individual control is
uncertain)

Small proportion
(individual
sensitivity)

**Under-nutrition,
Over-eating &
Malnutrition**

**Immediate or
progressive**

General or Groups

Burden of Foodborne Illnesses (USA)

**48 million Americans become sick due to foodborne illness each year
128,000 are hospitalized. Estimated annual deaths: 3000.**

(CDC estimations, 2010)

Pathogens that cause the most illnesses: Norovirus: 58%;
non-typhoidal *Salmonella* spp: 11%; *Clostridium perfringens*: 10%;
Campylobacter: 9%.

Leading causes of death: non-typhoidal *Salmonella* spp: 28%;
Toxoplasma gondii: 24%; *Listeria monocytogenes*: 19%; Norovirus: 11%.


Of 33,625 *Salmonella* isolates from animals & foods (BfR, 2000-08),
48% were resistant to at least one class of antibiotics and 35% were
resistant to more. Similar data for *Escherichia coli* & *Campylobacter*.

~203,000 visits/year to hospitals emergency depts for **food allergies**
(including ~90,000 cases of anaphylaxis).

Foodborne illness costs \$152 B/year (\$39 B due to fresh produce)

(“Health-Related Costs from Foodborne Illness in the United States”
by Robert L. Scharff, Produce Safety Project, 2010).

Obesity estimated at \$160 B/year in medical costs (Mc Kinsey Office)



Safety Hazards & Risks in the Food Supply Chain⁵

Drivers for Emerging Risks in Food Safety & Security

Overview (7 slides)

Driver 1: Globalisation

Growing global trade, migration & travel accelerate the spread of dangerous pathogens & contaminants in food.

Complex international supply chains or networks with long transit distances & time increase exposure to contaminants & complicate food safety management

Such supply chains are vulnerable to agro/bio-terrorist attacks.

One single contaminated food ingredient can lead to the recall of tons of food products in several countries, with high economic losses, possible import bans, damage to the tourist industry...

However, globalisation has many advantages:

Year-round supply of fresh food at competitive prices;
wide commercial opportunities with total annual exports of
agricultural & food products exceeding US\$ 1000 billion

Driver 2: Potential for Climate Changes

Water shortages cause **quantity & quality problems** with irrigation, process or ingredient water, + possible **shifts in production areas & cultured crops**, and **increased uses of agrochemicals**.

Flooding may cause increased **contamination of crops** in the field, or increased exposure of food animals to **zoonotic agents**.

Changes in temperatures & humidity are expected to affect the **distribution of plant & animal diseases**, the production of **mycotoxins**, the spread of food **pathogens**.

Changes in the oceans & coastal environments may affect **marine resources & seafood** (warming, algal toxins, acidification, pollution...)

FAO Monthly Food Price Index:

213.5 (June 2008); 206 (Nov. 2010); 214.7 (Dec. 2010)

Driver 3: Agriculture & Animal Breeding

Intensive agriculture & use of agro-chemicals will further impact on soil, water & energy supplies, with a high risk of **contamination** of water, food & feed crops **by fertilisers & pesticides**

Rising oil prices & the demand for agrofuels impact food **prices**

Animal breeding (FAO, 2010) **uses:**

~75% of world agricultural areas (including pastures & feed crops),
~8% of the world water consumption,
produces ~18% of total greenhouse gases.

The huge current increase in intensive production of eggs, milk & meat threatens the livelihoods of million of small breeders.

Large farms & herd size with enhanced contacts between animals, & their location near cities, **increase the risk of zoonotic diseases.**
3/4 of emerging human infectious diseases are due to animal pathogens
(SARS, BSE, avian Flu, Q fever, parasitic diseases...)

Antimicrobials given to animals as growth promoters & drugs contribute to **induce antimicrobial-resistant micro-organisms**

Driver 4: Trends in Food Processing

The trends for mildly preserved foods (minimal heating, cold pasteurisation, “natural” preservatives, “hurdle” technology) & for lower salt or sugar contents increase the risk of surviving pathogenic micro-organisms, possibly stress-resistant, & of microbial spoilage of foods.

Ready-to-eat foods with long refrigerated shelf-life enhance risks:

- Some manufacturers, especially SMEs, do not use appropriate product & process design, nor HACCP;
- Consumers do not always treat such foods as perishable items requiring refrigeration and/or specific conditions of preparation. Under-cooking is frequent.

Refrigerated storage has created niches for psychrotrophs: *Listeria*, *Yersinia*... & noroviruses.

Reduced packaging may also increase microbial risks & food wastes

Driver 5: Changes in Demographics

Global population is predicted to rise from 6.6 billion people in 2008 to 7.4-7.8 billion in 2020. The **world food demand** is further enhanced by increasing **affluent urban population and diet changes** in emerging countries (China, India...)

Many countries with insufficient production capacity will become **net food importers**.

Globally, due to climate changes & environmental constraints (water scarcity, risks of deforestation, loss of biodiversity...), **the main sustainable solution is to raise crop productivity on fertile land**

Overfishing has already decreased seafood resources

These trends will tend to **increase food prices** and reduce the global availability of certain foods.

Food choices will be affected,
depending on country & income level, towards foods requiring less land & resources (e.g. less beef, more poultry & pork)

Driver 6: Consumer Behaviour

Life expectancy & the **proportion of elderly people** will increase, with a higher **vulnerability to foodborne diseases**.

Changes in food composition motivated by consumer health
(less salt, sugar or anti-microbial agents)
may **enhance microbial growth**.

Increased **consumption of fresh, pre-packaged produce** or other food eaten **without heating** has already caused **pathogen outbreaks**.

Exotic food & ingredients, unusual animal species, small restaurants & catering, street vendors, also enhance risks & errors.

Organic food production & **animal welfare** (outdoor access) may lead to **reintroduction of pathogens with wildlife reservoirs** (*Trichinella*, *Toxoplasma*) or **increased prevalence** (*Campylobacter*).

WHO estimated the **number of overweight & obese persons**, to increase to 2,300 & 700 million in 2015, with **major health, social & economic impacts**.

In the EU: overweight persons = 50% of adults, including 15%¹²obese

Driver 7: Socio-Economic Changes

Cost is a major criterion for consumer's food selection, and manufacturers' **margins of profits** on food products are low, making it **difficult for firms to target investment to food safety**, & possibly enhancing fraud.


Poverty, city overpopulation, economic crises, increasing food prices, compromise food security, but also food safety and quality:

- **Food producers & processors may comply less with regulations** (use sick animals, recycle foods, do not recall contaminated foods, neglect factory maintenance...)
- **Low cost imports** (e.g. from aquaculture & seafood, fresh produce) may originate from regions with **poor sanitary practices**.
- Consumers may reduce the consumption of nutritious protein-rich animal foods, use foods past their expiry date.
People with reduced immunity will be particularly vulnerable

Social & economic inequalities, conflicts, migrants, natural disasters disrupt food security, and require large emergency operations (with complex logistics) that do not solve the development problems

Some Food Risk Management Tools

Risk Analysis



Risk = f (probability + severity) of adverse health impact of a hazard¹⁵

Quantitative Risk Assessment (QRA)

Proposals for Allergen Management in the Food Supply Chain

QRA implies the determination of “action levels”, i.e. maximum dose of an allergen present in a food portion which would not elicit severe reactions in the vast majority of allergic individuals

(for ex. ~ 8 mg of whole peanut elicit a **mild** reaction in only **10%** of the at-risk population).

With proper management of cross-contamination in the food industry (segregation, cleaning, traceability...),
3 categories of foods could then be offered:

- 1) “Free from...” (allergen not detectable)
- 2) “Suitable for...” (allergen below “action level”)
- 3) “May contain...” or “not suitable for...” (above “action level”)

Quantitative Microbial Risk Assessment

Risk pathway for *Salmonella* in chicken (T.P. Oscar, 2010)

Post-processing food handling unit operations	Associated pathogen events	Incidence (%) (data collection)
• Packaging (in the plant)	Low “initial” contamination level. Strain of high or lesser virulence. (= initial hazard identification by microbiological testing)	a
• Distribution • Washing • Cooking • Serving	Growth (if temperature and/or time abuse) Incomplete removal Survival (if undercooking) Cross-contamination	b c d e
Host demographics	Host vulnerability (aged, ill, on anti-acid medication) (response doses for given virulence strains)	f

————→ ASSESSMENT OF PATHOGEN EXPOSURE & RISK

Risk/Benefit Health Assessment of Food

For instance when the same food contains substances with adverse effects & substances with beneficial health effects.

The balance can shift towards benefit or risk depending on concentrations, intakes, subgroup of population.

Example: the potential benefit of eating fish & seafood in adults versus the potential risk of neuro-toxicity in infants and children due to methyl-mercury & persistent organic pollutants

When health risks do not clearly outweigh health benefits or vice-versa, the calculation becomes complex

Identification & Notification of Food Safety Risks

Rapid Alert Systems

Rapid Identification & Alert Systems

When food safety issues are first detected
(by public controls, food companies, consumers...)

they are usually related to specific hazard(s) and to given food chain(s).

**To make these data available to risk managers as soon as possible,
public authorities have created **rapid reactive systems**
for identification & notification of food hazards & health risks:**

- WHO International Food Safety Authorities Network (INFOSAN)**
- US Foodborne Diseases Active Surveillance Network (FoodNet)
(& US PulseNet for DNA fingerprinting)**
- EU Rapid Alert System for Food & Feed (RASFF)**

These are essential for risk & crisis management.
Their retrospective statistical analysis indicate trends,
but they are not predictive.

The EU Rapid Alert System for Food and Feed (RASFF)

lists food safety hazards (presence of illegal substances or unacceptable concentrations of chemicals or pathogenic organisms) **on its database, on a public weekly basis.**

Each Member State identifies hazards, in country or at EU borders, & must report them rapidly to RASFF, including measures taken regarding food safety (e.g. food recalls, rejection of not complying imports...)

RASSF sorts out the information and further transmits it to all M.S.


RASFF acts as a notification, communication & alert platform

RASFF has 34 members + 60 connected non-member countries.

<http://ec.europa.eu/rasff>

<https://webgate.ec.europa.eu/rasff-window/portal/index.cfm?event=notificationsList>

RASFF Annual Report 2009: Basis for Notification



~ 8000 notifications in 2009
including:

1796 “market notifications”
(578 “alert” + 1218 “information”)
(~ 50% from 3rd country products)
& 1484 “border rejections”²²

TABLE 1: ALERT NOTIFICATIONS

Notifications in blue typeface concern feed, all other notifications concern food.

DATE	NOTIFIED BY	REF.	REASON FOR NOTIFYING	NOTIFICATION BASIS	STATUS
30/06/2009	Italy	2009.0831	mercury (2.63 mg/kg - ppm) in smoked swordfish (<i>Xiphias gladius</i>) from Denmark	official control on the market	distribution on the market (possible) / product (to be) seized
30/06/2009	Hungary	2009.0833	deoxynivalenol (DON) (1464; 1613; 1398; 819; 2002; 1507; 2150; 1302 µg/kg - ppb) in durum groats from Hungary	company's own check	distribution on the market (possible) / product (to be) withdrawn from the market
02/07/2009	Poland	2009.0845	histamine (1433.5; 959.7 mg/kg - ppm) in frozen tuna steak from Spain	official control on the market	distribution on the market (possible) / product (to be) withdrawn from the market
02/07/2009	Germany	2009.0847	aflatoxins (B1 = 11.3; Tot. = 11.3 µg/kg - ppb) in groundnut kernels from Sudan, via the Netherlands	company's own check	distribution on the market (possible) / product (to be) returned to dispatcher
02/07/2009	France	2009.0848	Diarrhoeic Shellfish Poisoning (DSP) toxins (positive) in frozen cooked mussels without shell from Germany	official control on the market	distribution on the market (possible)
03/07/2009	France	2009.0849	<i>Escherichia coli</i> O157:H7 in frozen beef from Germany, via the Netherlands	company's own check	distribution on the market (possible) / product (to be) returned to dispatcher
03/07/2009	Austria	2009.0850	undeclared milk ingredient (casein: 1300 mg/kg - ppm) in dark chocolate bars from Italy	official control on the market	distribution on the market (possible) / product (to be) withdrawn from the market
03/07/2009	Austria	2009.0851	undeclared milk ingredient (casein: 810 mg/kg - ppm) in soybeans, roasted in the Netherlands, coated with dark chocolate manufactured in Belgium and packaged in Austria	official control on the market	distribution on the market (possible) / product (to be) withdrawn from the market
03/07/2009	Finland	2009.0854	norovirus (genogroup 2) in frozen raspberries from Poland	food poisoning	distribution on the market (possible) / product (to be) withdrawn from the market
03/07/2009	the Slovak Republic	2009.0858	undeclared nuts (134.10 mg/kg - ppm) in wafers with cocoa cream from Bulgaria	official control on the market	distribution on the market (possible) / product (to be) relabelled
03/07/2009	the Netherlands	2009.0860	benzo(a)pyrene (16.6 µg/kg - ppb) in food supplement from Germany	official control on the market	distribution on the market (possible) / product (to be) withdrawn from the market

RASFF Annual Report 2009

Alert Notifications

Risks most frequently reported from:

pathogenic micro-organisms *Listeria m.* (fish, dairy prod.)

Salmonella (feeds, meats & poultry...)

allergens (↑ undeclared milk ingredients or sulphites)

heavy metals (Hg in fish; Cd, in crustaceans & cephalopods)

mycotoxins (↓) (mainly in nuts, seeds, dried fruit & veg., spices).

55 recorded/notified cases of food poisoning, with 5 large outbreaks:

- Norovirus in frozen raspberries from Poland (2x)
 - Histamine in tuna from Sri Lanka
 - *Salmonella* from Spanish eggs
 - *Salmonella* from US peanut products

54 recorded cases of fraud (most for food of animal origin)

41 cases being from 3rd countries:

- Illegal imports • False health certificates • Unauthorised establishments

**An alert notification commands immediate action,
e.g. product withdrawal from the market**

RASFF Annual Report 2009

1484 EU Border Rejections

Products frequently rejected	Main rejection causes
1) Nuts & seeds	Mycotoxins
2) Fish & fishery products	Insufficient controls; parasites; heavy metals; microbiological hazards; biocontaminants
3) Fruits & vegetables	Mycotoxins, moulds, pesticides, pathogens, foreign bodies

**The control frequency of food imports is risk-based
& frequently adjusted**

Prediction (& Prevention) of Emerging Food Safety Risks

Predictive Early Warning Systems

“Expecting the unexpected”

**Assessment of emerging risks requires early detection or prediction
allowing preventive measures, before they become real risks**

Predictive early warning systems

Focused on specific food hazards:

Predictive microbiology

**Ex.: Early warning systems for molds & mycotoxins in maize or wheat:
Computer models based on weather variables**

Focused on food imports:

Predictive Risk-Based Evaluation for Import Compliance Targeting (PREDICT) automatically scans & screens many databases for information that may affect admissibility of food imports:

• a product's inherent risk rating (e.g., raw seafood); • if a company or product is subject to an import alert • the compliance history of relevant companies and product, including recalls, inspection results, analyses... • open source intelligence. PREDICT then issues a risk score for each entry line. Based on the score, the system may flag the entry as risky and send it to FDA

Vulnerability assessment:

CARVER-shock approach, for Food Defence against attacks

Vulnerable points in the food supply chain where the introduction of (intentional) hazards is most likely to occur.

shield the food supply against threats of terrorist attacks & new dangers. “Critical nodes” are identified, rather than specific hazards. Multidisciplinary experts focus on the production chain & flow diagram for a specific sector or product.

Predictive Early Warning Systems

Moderately focused: Data recording + expert or stakeholder opinion:

Sharing of information & knowledge to overcome problems of misinterpretation of data and ignorance of emerging risk signals.

EFSA's Stakeholder Consultative Platform, with web-based posting of documents, for feedback.

Broadly oriented: Combining food safety with other data:

Eurostat (EU Statistical Office): database “from farm to fork” which can show trends & developments associated with various steps in the production-trade-consumption chain in Europe.

FAO Global Information and Early Warning System on Food & Agriculture (GIEWS): preventive measures.

For ex.: **Satellite observations for agricultural land status & forecast crop harvest yields.**

Strategic foresight by experts and stakeholders:


Horizon scanning: systematic exploration of potential threats, opportunities & future developments

What policy changes are required to reach desirable scenarios?

EC Scientific Committee for emerging & newly identified health risks.

Safety prediction systems based on a holistic approach

Framework of the EU Project SAFE FOOD Structured Approach



Predictive systems based on a holistic approach

(after Marvin et al., 2009)

The EU project SAFE FOOD attempts to expand the scope of risk assessment & consider many sectors in & out the food production chain

science, technology, industry & innovation; international trade; nature & environment; government & policies; consumer behaviour; culture & demography; public health & welfare; agriculture; economy; information

Also attempts to identify specific indicators from these sectors:

Indicator = reliable & sensitive element informing on hazard & source of risk

Signal = change in such indicator, which may result in the development of a risk.

Multidisciplinary experts are asked to select critical indicators for a specific food supply chain and to score possible change in each indicator & subsequent consequences on food safety

Safety Risks in Food Production & Processing

Potential Safety Risks of Food Processing

Under-processing (including non homogeneity of treatment)

- Incomplete physical removal of contaminants
- **Incomplete inactivation of micro-organisms, spores, viruses, parasites**
(special risks for the chilled storage of foods of animal origin)
+ microbial recovery & adaptation of stress-resistant microbial strains
- Incomplete inactivation of natural toxicants, anti-nutrients, allergens

Over-processing (including non homogeneous treatment)

- Nutrient losses (& decrease in nutrient bioavailability)
- Formation of toxic molecules (acrylamide, heterocyclic amines, HMF..)
- Enhancement of microbial growth (slicing, grinding, long storage...)
- Migration of chemicals from package into food (including inks...)

Intentional addition of unsafe substances


some food additives, nanoparticles, adulteration products,
antibiotic-resistance genes, pathogens...

Acrylamide in Foods (discovered in 2002)

Acrylamide may be formed in carbohydrate-rich foods: potatoes & cereal-based products (chips, French fries, bread, crispbreads, biscuits, breakfast cereals, coffee) during cooking processes such as **frying, baking, roasting, microwaving at $\geq 120^{\circ}\text{C}$ (not boiling)** (up to 5 mg/kg potato chips and 2 mg/kg biscuits).

It may also form in heated (oxidised?) fat-rich dry foods such as nuts.

Exact mechanisms of formation are unsure, but involve Maillard reactions between asparagine & reducing sugars/reactive carbonyls



prop-2-enamide ($\text{C}_3\text{H}_5\text{NO}$)

Acrylamide is also a highly reactive industrial chemical used for the production of plastics & materials.

Acrylamide is a genotoxic carcinogen to experimental animals, EU listed as Substance of Very High Concern, thus exposure from all sources, including diet, should be minimized. Its primary metabolite, glycidamide, is also toxic.

Reducing Acrylamide Formation

CIAA “Toolbox” to show how recipes & processes can be modified (use low sugar varieties & mixes, avoid overheating & brown colour, add asparaginase or amino acids or an asparagine-degrading yeast).

From 2003 to 06, the acrylamide levels of EU bread & potato chips has decreased, but not that of biscuits, breakfast cereals, French fries, cereal-based baby foods (EFSA).

The link between acrylamide & cancer risk in humans is uncertain. People have been exposed to acrylamide in their diet for a long time.

Recent studies suggest that tolerable daily intakes of acrylamide for carcinogenic levels should be set at 2.6 µg/kg b.w., while the TDI for neurotoxicity is higher, at 40 µg/kg b.w.

Canadian, Swedish & US studies estimate the average exposure of adults to acrylamide in food as 0.3, 0.5 & 0.4 µg/kg b.w. per day, respectively.

This example shows how even long-used processes can suddenly reveal unsuspected risks (frying & grilling of meat & fish may form carcinogenic heterocyclic amines. HMF, furan, acetaldehyde -potential carcinogens- may also be formed in various foods upon heating)

“Margins of Exposure” (MoEs) to Carcinogens in Foods

Attempt to quantify risks by calculating the ratio:

Dose that produces a certain frequency of cancer in animals/
divided by the **dose that humans are exposed to.**

Low value MoEs are probably dangerous

Genotoxic carcinogens	MoE (JEFCA, 2005) (for median human intakes)
Acrylamide	300
Ethyl carbamate	20,000
Polycyclic aromatic hydrocarbons (PAHs)	25,000

Although the risk from acrylamide is much larger than that from the 2 other (regulated) carcinogens (but less than from smoking), there are no regulations concerning acrylamide in foods.

Hormonal Disruptors from Food Packaging & Containers

Chemicals used as plasticizers in food packaging and various plastics, such as **phthalates, bisphenol A... are suspected of causing increasing numbers of anomalies.**

Bisphenol A (BPA) is used in polycarbonate bottles, infant feeding bottles & in epoxy resins serving as protective coatings for food & drink cans & vats

During storage, BPA migrates in small amounts into food & drinks.

EFSA (2007) indicates that the intake is 5 to 500 fold below the “tolerable daily intake”, and that BPA is metabolised & eliminated.

BPA is permitted in food contact materials in the EU, USA, Japan.

Canada and the EU have banned its use for infant food bottles (2010)

Frauds & Adulterated Foods

Profit-motivated frauds, often criminal, have caused many food crises (e.g. direct or indirect addition of toxic substances to foods or feeds):

Examples	Impact	Fraud (& error)
Spanish oil syndrome (1981)	20,000 persons affected >800 deaths	Denatured oil sold as olive oil
Ukrainian sunflower oil (2008)	Exported to EU Late withdrawal & ban	Mineral oil added
Bovine spongiform encephalopathy (UK, 1984, then EU)	>100 deaths >1 million cows culled ↘ meat consumption Huge economic losses Loss of consumer trust Changes in regulations	Prion-infected meat & bone flours fed to cows. Milder flour process. Ban on flours not respected
Belgian chicken (1999)	Economic losses	Dioxin in feeds (from electric transformer oil)

Frauds & Adulterated Foods (2)

Examples	Impact	Fraud (& error)
Irish pigs & pork meat products (2008)	Large trace & recall from 54 countries	PCB & dioxin in feeds 100 fold legal limit. Dried/contaminated fuel
German feeds, pigs, chicken, eggs (2010)	~150,000 T feeds, 6000 farms affected	Dioxin-rich industrial fatty acids added to feeds
Chinese gluten, milk, infant milk formulae (2007-2010)	Grave kidney problems for >300,000 infants & children	Melamine addition to fake higher N content + long denial from authorities

Non authorised food or feed additives are frequently used in some countries (with corruption, official silence, opaque supply & export channels).

In 2007, some 2000 Chinese producers & 1000 retailers had exported food ingredients & additives worth US\$ 4 billion (including vitamins & amino acids); Food & pharmaceutical companies worldwide used these ingredients & additives without sufficient traceability or quality control.

Counterfeited Foods & Drinks

The global level of fraud as counterfeiting in the food & drink industry is estimated at ~ \$50 bn/year.

Global trade & high food prices increase fraud opportunities

High losses for food firms through damage to brands & spending on security measures. Also **health risks for consumers.**

Security measures on packaging include: holograms, microdots, data codes, use-by dates, lot numbers, markers... **are not always efficient**

The EU plus 10 countries (Japan, USA, Korea, New Zealand, Canada, Australia, Switzerland, Morocco, Mexico, Singapore) have been negotiating an **anti-counterfeiting trade agreement (ACTA)**

Counterfeited Foods & Drinks



**Destruction of faked whiskies & cognacs
at Zhuliao (Guangdong province)**

Potential Food, Feed or Drink Targets for Bioterrorism

(intentional addition of toxic chemicals or biological agents)

Those with easy access, large volumes, wide diffusion, short shelf-life, able to reach a maximum of people, or a specific group of people

- **Water:** rivers, ponds, city reservoirs, tanks, sea salt ponds
(although protected by water treatments, controls & dilution effect)

- **Liquid foods:** milk, fruit juices, beer, wine...
Liquids facilitate an uniform diffusion of contaminants

- **Ingredients** (possibly “foreign”) which are sourced globally, at low prices, undergo less controls, are less visible and reach multiple end-products: flours, milk powder, salt, sugar, tomato paste, peanut butter..

- **Counterfeited foods with faked known & trusted brands:**
Coca Cola, Nescafé, baby foods brands...

- **Food Animals** (& animal feeds), which can transmit zoonotic diseases to man through consumption of animal products.

Safety of “Innovative” Food Technologies

“New” Preservation Technologies: Potential Food Safety Risks

Physical treatments (often minimal, or low temperature)

High pressure pasteurisation: incomplete microbial inactivation, no spore inactivation, microbial recovery; some chemical reactions

Pulsed electric fields: no spore inactivation, non homogeneity & arcing; electrochemical reactions; metal transfer from electrodes?

Cold plasma (dielectric barrier discharge at atmospheric pressure and $\sim 30^{\circ}\text{C}$). OK for surface disinfection, but free radicals, oxidations?

Light pulses & UV: non homogeneity (surface only); photo-oxidative reactions.

Ultrasounds: safe cleaning but low efficiency/microbial inactivation

Ohmic heating: under/overheating, metal transfer from electrodes?

Microwaves: non homogeneity (cold spots) & power reduction over time of domestic MW ovens?

Ionising radiations: free radicals, oxidative reactions, consumer distrust (US petition for surface irradiation of meat carcasses)

Chemical or combined treatments

Modified atmospheres, anti-microbial agents, “hurdle” processing (often minimal or low temperature processing)

- Incomplete microbial inactivation
 - Growth only delayed in storage
 - Inciting to over-extend shelf-life
- Sub-lethally injured cells may give antimicrobial-resistant, acid-resistant, osmotic-resistant or virulent strains, through adaptation.
 - Potential reactivity or toxicity of antimicrobial agents
 - Possible use to conceal poor hygiene practices

Decontamination of poultry, pork or beef carcasses by washing/spraying with solutions of Cl_2 , ClO_2 , hypochlorite, acidified Na chlorite, trisodium phosphate, organic acids (lactic), H_2O_2 , O_3 , electrolysed H_2O , lactoferrin...
Only some of these antimicrobials are authorised in the EU.

Fresh-cut bagged fruit & vegetable: Norovirus, *Salmonella*, *E. coli* O157 (18 major outbreaks in the US 1998-2006, mostly due to leafy greens).

Initial contamination through irrigation water & fertilizers.

Increased surfaces from cut or peeled tissues

release liquids & nutrients, enhance microbial attachment & growth, and interfere with disinfectant washes (NaOCl , Cl_2) or CO_2 M.A.

Nanotechnologies

Use of engineered substances on a very small scale: 1-100 nm (obtained by assembly of molecules or downsizing).

Nanoparticles have specific physico-chemical properties:
Small size increases diffusivity. High surface area ($>60 \text{ m}^2/\text{cm}^3$)
enhances reactivity, binding, recognition...

Structure, size distribution, chemical composition, & surface charge may also confer **unique functionalities for food uses:**

1) As food ingredients & additives: increased solubility, dispersion; stable emulsions without emulsifier; low fat emulsions

2) As encapsulation & delivery systems for bioactive compounds: molecular traps (e.g. emulsion droplets) for protection & targeted delivery (lycopene, phytosterol); **higher nutrient availability** (Fe, Zn)

3) As more protective or intelligent packaging: in polymer or biopolymer films & coatings, nanoparticles act as “fillers” to enhance **barrier properties & resistance to mechanical or thermal stress.**

Reactive nanoprobables alerting of use-by date, temperature, pathogens, toxins;
Anti-bacterial coating (silver particles) for food-contact surface
(refrigerator)

Safety of using nanoparticles in foods? & health implications of exposure

(EFSA Draft Opinion, 2008)

Existing risk assessment methods can be applied, **on a case/case basis**

Data on non nano chemicals **cannot be extrapolated** to their nano equivalents, **since formulation to the nanosize may change properties**

There are considerable limitations & uncertainties: 1) on **detecting, characterising & “dosing” ENM**; 2) on their **absorption, distribution** (in fatty tissues, bone, brain?), **accumulation, metabolism & excretion**; 3) on their **toxicity**

Small size increases their ability to move unexpectedly in the body.
Adhesive & reactive surfaces may cause various interactions.

Main other challenges (many other uses of nanoparticles):

workplace safety; distinction of natural & engineered nanoparticles; cost; impact on the environment: biodegradation/persistence/accumulation in soil, water, air, food chain; consumer uncertainty & suspicion; labelling & legislation.

Active & Intelligent Food Packaging: Safety Concerns

While “classic” packaging should be as inert as possible, **these new packaging intentionally interact with the food or its environment**,
1) either (“active) to extend shelf-life with maintenance of quality,
or 2) (“intelligent”) to give indication & monitor the food freshness
(time-temperature indicators, ripeness indicator, biosensors, tamper-proof or identity-ensuring systems, nano-chips in smart inks...).

Active packaging contain deliberately incorporated components intended to release or absorb substances into or from the food or its environment (release antimicrobials, antioxidants; absorb O₂, C₂H₄)

Highly promising, these new packaging are not yet widely used

The main safety issue (as for classic packaging) is migration of chemicals (& their degradation products) into the food.

Nano sizes could increase risks.

A recent Regulation (450/2009/EC) specific to A&I packaging introduces safety evaluation by EFSA, with an authorisation scheme, focusing on migration data of chemicals & their toxicological properties⁴⁷

Active & Intelligent Food Packaging: Safety Concerns (2)

Their efficiency to perform the claimed function should be demonstrated in real foods. This is critical when they should prevent microbial growth or reveal the presence of pathogenic bacteria or toxic contaminants.

Their proper use should be explained to consumers by way of labelling.

Their acceptance may be limited since Consumers may perceive systems for the extension of shelf-life as detrimental to food freshness, while **Retailers** fear that they may induce consumers to select only recently displayed items, and convey a negative carbon footprint image.

RFID tags printed on food packages (no direct interaction with food) may change food sales & marketing, allow cell phones to indicate food composition, nutritional adequacy; refrigerators to signal expiry dates & send automatic reorders...

with some risks of “neuromarketing” & intrusion into private life!

Citizens' Perception of Risks in Food Technologies

Technologies which tend to elicit consumer suspicion & rejection
(& strong political debates):

- Ionising irradiation of foods
- Hormonal (& antibiotic) treatment of animals to hasten growth & increase meat or milk production (banned in the EU)
- Genetically modified food crops & food ingredients (in the EU)
- Genetically modified animals (cloned animals & their offspring)
- Various food additives (consumers' request for “clean labels”)

The **consumer right to informed choice** is well established in the EU:


Mandatory labelling for irradiated foods and GM foods have discouraged manufacturers & retailers to place such foods on the market, in spite of their potential advantages.

Survey on How European Citizens Perceive Food-Related Risks

EFSA 2010 - Eurobarometer 354 - TNS Opinion & Social, Brussels

Face to face interviews with 27,000 individuals >15, in 27 M.S.

Spontaneous responses



Negative attitude to “synthetic” chemicals, but insensitivity to dose-response relationship. Preference for “natural” processes.

Eurobarometer Survey 354 (EFSA, 2010) (2)

- **Trust in information sources:**

Doctors & health professionals: 84%	Family & friends: 82%
Consumer organisations: 76%	Scientists: 73%
Environmental protection groups 71%	Food safety agencies: 64%
European Institutions 57%	National governments 47%

- **The role of public authorities is perceived as effective, but**
1 in 2 consider that they are not independent from producers' interests.

According to the EU Commissioner for Health,
conditions necessary to support acceptance of innovation are:

- 1) **Clear benefits perceived for consumers & society**
- 2) **Predictable regulatory environment & legal certainty**
(+ safety assurance & control)
- 3) **Science behind risk assessment made more credible**
- 4) **Improved transparency & dialogue with the public**

Thank you for your attention

Future EU Regulation on Novel Foods & Novel Ingredients

Disagreements and delays

The **E. Council** wants “**nanofoods**” included. The **E. Parliament** wants a moratorium until safety is established, and asks for indicating any nanomaterial on the food label.

The **Council** supports including food from the **offspring of cloned animals**, while **Parliament** wants to ban such food, and asks for a new law covering ethical & safety issues of animal cloning.

The **E. Commission** proposed (Oct. 2010) a 5 year ban on the use of cloned farm animals and on the marketing of food from cloned offspring, although it supports **EFSA**’s opinion on the lack of food safety problems.

The **E. Commission** recognises the problems of animal welfare and ethics.

ENDORSED FOR PUBLIC CONSULTATION

DRAFT SCIENTIFIC OPINION

Guidance on risk assessment concerning potential risks arising from applications of nanoscience and nanotechnologies to food and feed¹

EFSA Scientific Committee^{2, 3}

January 2011

www.efsa.europa.eu/en/consultations/call/scaf110114.pdf

The EU Food Legislation on Novel Foods & Ingredients (1)

Protection against Food Safety Risks or Barrier to Innovation & Trade?

Regulation 97/258/EC on Novel Foods & Food Ingredients subjects each **novel food to a severe pre-market safety assessment & authorisation procedure**. Also edicts **specific labelling rules to inform the consumer** of any characteristics making the novel food no longer “equivalent” to an existing food, or having health or ethical implications.

Categories of Novel Foods:

- 1) a food not used significantly for humans in the EU before 1997**
- 2) a new or modified molecular structure or ingredient** isolated from animals, plants, micro-organisms, fungi, algae
- 3) a food or ingredient subjected to a new process, or issued from a new production or breeding process**, resulting in significant changes in composition, structure, nutritive value, metabolic effect and/or level of undesirable substances.

Includes “**functional foods**”, also subject to an **Health Claim Regulation**

Excluded from the Regulation (because other rules apply): food additives, flavourings, enzymes, extraction solvents, vitamins & minerals, GM foods.

Future EU Regulation on Novel Foods & Novel Ingredients

(2)

Regulation 97/258/EC on Novel Foods & Food Ingredients exerts strong detrimental constraints on innovation & trade (cost, delays), as much as a protection against food safety risks.

A revised proposal (of 14 Jan. 2008) is discussed, to:

- 1) Promote a more favourable legislative environment for technological innovation & industrial competition
- 2) Consider the particular needs of traditional foods from 3rd countries (a “notification procedure” may be sufficient)
- 3) Clarify & facilitate implementation, with a single EU harmonised, centralised & shorter procedure for assessment & authorisation
- 4) Give a wider choice of safe novel foods to the consumers.

Food Safety Management Tools

- Food legislation & standards
- GAP, GHP, GMP & HACCP certifications
- Food quality & safety management systems (ISO 9001 & 22000)
- Risk analysis

Others:

- Traceability
- Food operators' responsibility & auto-controls
 - Third party certification
- National food inspections & controls, including at borders
 - Risk-based controls
- Mandatory transparency & reporting of food incidents
 - Mandatory collaboration with authorities
 - Withdrawal & recall procedures
 - Prior notice of imported food shipments
 - Registration of food facilities
- Food defence plans of individual food companies

Selection of critical factors for identifying emerging food safety risks in dynamic food production chains

(Van Asselt et al., Food Control, 2010, 21, 919-926)

Endogenous indicators

Factors 1-6 relate to the **Food chain complexity**: the higher possibility of errors in complex systems results in food safety risks.

Factors 7-10 are linked to the **Producers' attitude towards food safety**.

Factors 11-13 show **Producers' compliance to food safety regulations**.

Factors 14-18 correspond to **Technical innovations** in the food chain.

Exogenous indicators


Factors 19-22 include **Origin of raw materials; Legal requirements; Climate change; & Economic status**.

Factors 23-29 are related to **Consumers: Size of demand; Demand for convenience or quality; Consumer concern for health**

Multidisciplinary experts were asked to select critical indicators for each specific case, rank them (individually) and detect & analyse their signals.

RASFF 2008

(POTENTIALLY) PATHOGENIC MICRO-ORGANISMS



	Classification	Date of case	Last change	Reference	Country	Subject	Product Category
1.	alert	20/12/2010	07/01/2011	2010.1725	FR	suspicion of phytohaemagglutinin in chilli con carne from France	prepared dishes and snacks
2.	alert	28/12/2010	07/01/2011	2010.1769	IT	parasitic infestation with Anisakis of anglerfish (Lophius piscatorius) from the United Kingdom	fish and fish products
3.	alert	28/12/2010	07/01/2011	2010.1771	DE	dioxins (in fat used for the feed: 123; 124 pg WHO TEQ/g) in vegetable feed fat for pigs and poultry from Germany	feed materials
4.	information	06/01/2011	06/01/2011	2011.0024	HU	unauthorised substance pseudovardenafil (suspected) in food supplement from Malaysia	dietetic foods, food supplements, fortified foods
5.	information	06/01/2011	06/01/2011	2011.0026	PT	prohibited substance nitrofuran (metabolite) nitrofurazone (SEM) (3.3 µg/kg - ppb) in frozen beef from Brazil	meat and meat products (other than poultry)
6.	alert	26/11/2010	06/01/2011	2010.1624	NL	hepatitis A virus (present) in dates from Algeria, via Belgium	fruit and vegetables
7.	border rejection	23/12/2010	06/01/2011	2010.CIC	GB	aflatoxins (B1 = 5.0; Tot. = 5.9 µg/kg - ppb) in peanut butter from Sierra Leone	nuts, nut products and seeds
8.	alert	06/01/2011	06/01/2011	2011.0025	DK	deoxynivalenol (DON) (1000 µg/kg - ppb) in organic oat meal from Denmark, with raw material from Sweden	cereals and bakery products
9.	alert	06/01/2011	06/01/2011	2011.0023	IE	undeclared soya in white bread flour from the United Kingdom	cereals and bakery products
10.	border rejection	06/01/2011	06/01/2011	2011.AAP	NL	aflatoxins (B1 = 16.6; Tot. = 18.8 / B1 = 5.7; Tot. = 6.4 µg/kg - ppb) in groundnut kernels from Argentina	nuts, nut products and seeds
11.	alert	23/12/2010	06/01/2011	2010.1752	IT	mercury (1.7 mg/kg - ppm) in swordfish loins from Spain	fish and fish products
12.	alert	06/01/2011	06/01/2011	2011.0022	IE	undeclared soya in batter mix from the United Kingdom Salmonella Derby (3 out of 11 samples /25g), Salmonella enteritidis (FT1 + FT14b: 7 out of	cereals and bakery products

Risk/Benefit Health Assessment of Food

J. Hoekstra et al., Food & Chemical Toxicology, 2010, in press




Fig. 2. A schematic description of the steps within each tier.




CERPTA



Necesidades en Tecnología Alimentaria

- Conservación de alimentos
 - Destrucción patógenos
 - Destrucción otros microorganismos alterantes
 - Inactivación de enzimas
 - No toxicidad
 - Compatibilidad de los envases con el alimento y el proceso
 - Nuevas emulsiones, nuevas texturas
 - Sistemas de control-uniformidad

Equipos de alta presión hidrostática





Economical analysis for novel process technology: High Pressure Processing

NC Hyperbaric
Carole Tonello, PhD

NovelQ – Sept 6, 2007

Instalaciones


Línea de alta presión



Línea de alta presión
homogenización



Ultra alta presión homogenización





Ultra alta presión homogenización

- ✦ Alta presión
- ✦ Impacto
- ✦ Cizalla
- ✦ Cavitación
- ✦ Turbulencia

Stansted Fluid Power

- ✦ Redesigned ceramic valve, changing its geometry
- ✦ On 92-340 MPa treatments flow speed reach 75-100 m/s before exiting the valve and 120-200 m/s after
- ✦ It's possible withstand up to 400 MPa




Patente EU solicitada UAB

- ✦ CONTINUOUS SYSTEM AND PROCEDURE OF STERILIZATION AND PHYSICAL STABILIZATION OF PUMPABLE FLUIDS BY MEANS OF ULTRA-HIGH PRESSURE HOMOGENIZATION
- ✦ Pat n° 10380094.2-1221
- ✦ Explotación. EBT YPSICON

Nuevo equipo UHPH 400 MPa 125 l/h. CERPTA-UAB (Pat n° 10380094.2-1221)



DISEÑO Y PRODUCCIÓN DE ALIMENTOS FUNCIONALES



Diferencia alimento funcional-medicamento

- ✦ El medicamento cura o cronifica una enfermedad
- ✦ El alimento funcional tomado regularmente puede reducir el riesgo de tener una determinada enfermedad

Tecnofuncionalidad

- ✦ Aplicación de procesos tecnológicos incluidas las nuevas tecnologías para mantener o mejorar la funcionalidad de un alimento, por las modificaciones que se producen a nivel conformacional o estructural

Tipos de alimentos funcionales

✦ Origen natural

✦ Alimentos funcionales de diseño




BIOCAT




BIOCAT

The Scale of Objects

Natural Objects



Manufactured Objects




Weiss, IFT. 2007



Productos en función del tamaño de la protección

- ✦ Microproductos: Cápsulas entre 1 y 3 micras
- ✦ Submicroproductos: Submicrocápsulas entre 100 y 1000 nm
- ✦ Nanoproductos: Nanocápsulas inferiores a 100 nm (Engineering Nanomaterials-EN)

Liposomas



Nanoencapsulación



UHPH hasta
400 Mpa







Fig. 1. Schematic view of the Stated high pressure homogenising valve.

ImagenTEM de submicrocapsulas






CERPTA

GRACIAS POR SU ATENCIÓN





SISTEMAS ACUOSOS – I

AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

www.malta-consolider.com



Matter at High Pressure



Grupo: Centro de Astrobiología

Felipe Gómez
Laura Jiménez Bonales
José Antonio R. Manfredi
Victoria Muñoz Iglesias
Olga Prieto Ballesteros

I. AGUA Y SISTEMAS RELACIONADOS CON LA VIDA

- a. Síntesis, propiedades y estabilidad de clatratos**
- b. Efectos de la alta presión en soluciones acuosas de agregados supramoleculares y proteínas**
- c. Microbiología en condiciones extremas de presión y temperatura**

Clatratos en Europa


Master de tesis de Vicky – Octubre 2010
“Clatratos en Europa, satélite de Júpiter”

En el sistema solar exterior se dan las condiciones y los materiales necesarios para que se formen clatratos de gas, tanto en los satélites de hielo como en los cometas

- agua
- volátiles
- P y T

Manifestaciones geológicas observadas en los satélites de hielo pueden ser producto de la descomposición de los clatratos de gas

- estructuras de terrenos caóticos
- actividad de géiseres



EUROPA

Europa es un satélite especialmente interesante para Astrobiología debido a que parece albergar en su interior un océano de agua líquida en contacto con un substrato rocoso.

Actualmente Europa tiene una superficie de hielo dominada por H_2O , sales hidratadas y/o hidratos de ácido sulfúrico, CO_2 , SO_2 y H_2S .

El océano de agua líquida está en contacto con la capa rocosa probablemente afectada por actividad hidrotermal, cuyas fumarolas liberan gases.



Estabilidad de los clatratos de gases volcánicos en el interior del satélite. De oxidantes radiogénicos en la subsuperficie.

APLICACIONES:

Terrenos caóticos

Diferenciación criomagmática

ESTABILIDAD CLATRATOS de CO_2



$\text{CO}_2 \rightarrow$ Estabilidad desde 0.05-0.5 mm de profundidad hasta el océano




CLATRATOS VS. SALMUERAS





Las áreas superficiales no dominadas por el hielo de agua, son ricas en hidratos de sales y de ácido sulfúrico.

Se cree que estos materiales tienen origen endógeno por su relación estructural con las fracturas y la presencia de un campo magnético autoinducido.

Tradicionalmente los electrolitos en solución se han descrito como inhibidores de formación de clatratos debido a:

- La fuerte atracción del agua hacia los electrolitos.
- El efecto “*salting-out*”: descenso de la solubilidad del gas en la salmuera debido al enlace del agua con los iones.










DIFERENCIACIÓN POR CLATRATIZACIÓN

La formación de clatratos desde un magma enriquecido en gases e iones disueltos puede producir la diferenciación de la cámara magmática en los satélites de hielo.

Durante el proceso, las sales disueltas pueden precipitar y separarse por su mayor densidad, permitiendo a la solución con gases cristalizar en clatratos de gas.

Si los clatratos se desestabilizan por descompresión, los criomagmas ricos en volátiles podrán ascender a la superficie, e incluso erupcionar explosivamente.



CO₂ en disoluciones acuosas planetarias



(desarrollo de Laura)

- CO₂ en soluciones sulfatadas a alta presión

Aplicación en modelos de criomagmatismo del SSE. Explicación de la flotabilidad de los criomagmas. Explosividad de las erupciones de criomagmas. Diferenciación por clatratación

- CO₂ y carbonatos

Aplicación en la formación de carbonatos en épocas tempranas de la Tierra y Marte




Carbonatos en Tierra y Marte primitivos

Los carbonatos en la Tierra desempeñan un papel muy importante en el ciclo del C, fijan y liberan CO₂

Pueden ser precipitados inorgánicamente o ser biomedidos

Actualmente los carbonatos precipitan en mares someros, según la reacción:



Los carbonatos son geomarcadores de unas condiciones ambientales particulares. Y pueden ser biomarcadores



Existen periodos en la historia de la Tierra con carbonatos particulares **¿Cómo precipitan?**

- Carbonatos arcaicos
- Carbonatos proterozoicos


¿Dónde están los carbonatos en Marte?

¿Se puede precipitar la dolomita inorgánicamente en condiciones de la corteza terrestre?

635 Ma cap-carbonate sequence, NW Namibia



Tierra primitiva-carbonatos vs. Alta P




Proterozoico- los “*cap carbonates*” son calizas y dolomías que aparecen por encima de los depósitos glaciares proterozoicos (Makganyene ~ 2220 M.a., Sturtian ~700 M.a. o Marinoan 635 M.a.). Tienen características especiales en relación a carbonatos de otras épocas: cementos de aragonito bien desarrollados, ondulaciones gigantes, barita diagenética, relación isotópica deficitaria en ^{13}C etc

Los carbonatos del Arcaico (más de 2400 m.a.) no son frecuentes en la Tierra. Tienen características peculiares:

- Asociaciones de caliza y ortocuarcitas (sedimentarias o volcánicas)
- Carbonatos asociados a esquistos o cherts
- Carbonatos en asociaciones volcánicas
- Sedimentos clásticos-carbonatados

Arcaico-atmósfera más densa. Hasta 200 bares de presión.





Carbonatos Proterozoicos. *Cap carbonates*

Carbonatos que están en contacto limpio con facies periglaciares. Son dolomías microcristalinas que aparecen por todo el globo, con calizas post-glaciares por encima


Se han propuesto 3 modelos que relacionan una deglaciación abrupta y el evento de precipitación de carbonatos (alcalinidad extrema)

- **cambio brusco de los océanos** anóxicos profundos, desgasificación de CO_2 , subsecuente liberación de la alcalinidad en nuevas cuencas subsumergidas y precipitación de CaCO_3
- **Snow Ball Earth**: alto grado de alteración química debido a condiciones de un super efecto invernadero y de una glaciación global. Deglaciación catastrófica cuando el nivel de pCO_2 fue crítico (0.12 bares)
- liberación masiva de la alcalinidad por desestabilización de **clatratos de metano**, debido a la oxidación anaeróbica por sulfato-reducción



Experimentos para la posible comprobación de las hipótesis de SBE y los clatratos (baja temperatura, despresurización, precipitación de dolomita???)

Because of an extended cold spell, oceans start freezing.



Lowered reflectivity causes further cooling, ending in "snowball Earth."




CO₂ cycle in ocean stops; CO₂ outgassed by volcanoes builds up.





Strong greenhouse effect melts "snowball Earth," results in "hothouse Earth."



CO₂ cycle restarts, pulling CO₂ back into oceans, reducing greenhouse effect to normal.



THE CARBON CYCLE



[Processes lettered in blue are absent in a snowball Earth]



Carbonatos Arcaicos

La atmósfera del Arcaico fue muy rica en CO_2 . Esto pudo acidificar los océanos. Condiciones no favorables.

Los datos muestran que la alteración de las rocas ígneas liberó Ca^{2+} a los océanos

La corteza era delgada e inestable, con pocos lugares aptos para la precipitación masiva de carbonatos. Solo en partes profundas con favorable contenido en CO_2 (fáciles de destruir posteriormente por tectónica)

¿Cómo precipitaron los carbonatos del Arcaico teniendo en cuenta las características de la atmósfera y del océano? Desde pCO_2 alta y T medias





Marte primitivo-carbonatos vs. Alta P

Si Marte tuvo agua y una atmósfera mas densa podría haber formado carbonatos

El rover Spirit ha identificado carbonatos Mg-Fe (16 - 34 peso%) en Columbia Hills del cráter Gusev. Podrían haber precipitado de soluciones carbonatadas en condiciones hidrotermales a pH casi neutro en asociación con la actividad volcánica del periodo Noico

¿Son hidrotermales los carbonatos marcianos?



CO₂ EN DISOLUCIONES ACUOSAS PLANETARIAS

www.malta-consolider.com



Matter at High Pressure

PROGRAMA
ingenio
2010




CAB
ASOCIADO AL NASA ASTROBIOLOGY INSTITUTE

CARBONATOS Y SU CLASIFICACIÓN

CALCITA


Calcita	CaCO_3
Magnesita	MgCO_3
Siderita	FeCO_3
Rodocrosita	MnCO_3
Smithsonita	ZnCO_3



Romboédrico

DOLOMITA


Dolomita	$\text{MgCa}(\text{CO}_3)_2$
Ankerita	$\text{FeCa}(\text{CO}_3)_2$
Kutnahorita	$\text{MnCa}(\text{CO}_3)_2$



Romboédrico

ARAGONITO

Aragonito	CaCO_3
Witherita	BaCO_3
Estroncianita	SrCO_3
Cerusita	PbCO_3



Rómbico


CARBONATOS SÍNTETICOS Y ANÁLISIS

✓ Síntesis de carbonatos en C.N.


✓ Comportamiento de carbonatos
A distinta P y T una vez formados

✓ Difracción de RX
✓ Espectroscopía Raman
✓ SEM y TEM
✓ Modelización geoquímica
de disoluciones
✓ ICP-Masas de disolución y
sólidos
✓ Isótopos estables (C,O)

✓ Síntesis
a 25°C y 1 bar



✓ Síntesis
a 25°C y 1 bar



CARBONATOS SÍNTETICOS:

OBJETIVOS

1º Síntesis en condiciones del Proterozoico ($\uparrow P$, $\downarrow T$)

2º Síntesis en condiciones del Arcaico ($\uparrow P$, $\downarrow T$)





- ✓ Análisis “in situ” de las reacciones
- ✓ Análisis “ex situ” de los minerales



CARBONATOS SÍNTETICOS

✓ Síntesis en condiciones
del Proterozoico:


1. A partir de HCO_3^- y H_2CO_3
2. Bajas temperaturas (0°C y -5°C)
3. Presiones desde 6-100 bar



METODOLOGÍA: Síntesis de CaCO_3 (Proterozoico)



Experimentos de compresión y descompresión a bajas T



METODOLOGÍA: Ejemplo, Síntesis de CaCO_3 (Proterozoico)



✓ Síntesis
a 25°C y 1 bar
(estudio previo)

Desplazamiento de la reacción
(función del pH y concentración),
Cinética rápida
Rendimiento al 100%





✓ Síntesis
Proterozoico:
a bajas T y altas P

Desplazamiento de la reacción
(función del CO_2),
Cinética lenta
Rendimiento % < 100%, restos de HCO_3^-

ANALALISIS “in situ”

seguimiento mediante espectroscopía




Raman de la disminución de la
concentración de HCO_3^-




Estudio del HCO_3^-



pH




ANALALISIS “ex situ”




RAMAN

RX







CARBONATOS SÍNTETICOS



- ✓ Síntesis en condiciones del Arcaico:
1. A partir CO_2
 2. Temperaturas medias (20°C)
 3. Presiones hasta 200 bar


METODOLOGÍA: Síntesis de CaCO_3 (Aracaico)



Estudio de la solubilidad del CO_2 en función de la p , T y presencia de sales

1. Carbonatos en la tierra primitiva

2. Clatratos: $\text{CO}_2 + \text{H}_2\text{O}$



Estudios cuantitativos
mediante
Espectroscopía Raman

Metodología



Ej: MgSO_4 en H_2O

Estudios cuantitativos
mediante
espectroscopía Raman:


$$I = N_i \sigma(\nu_{ex}) \nu_{ex}^4 E_0 e^{-E_i / kT}$$



Intensidades o
Áreas relativas
Varía en función de:
1. P
2. T
3 concentración de sales





3325 cm⁻¹




Calibrado

Calibración para CO₂

- CO₂(g) → CO₂(ac) vs. P, T en H₂O



Solubilidad del CO₂(g) en disoluciones



Carbonatos en
tierra primitiva

Clatratos


GRACIAS POR SU ATENCIÓN



Muerte y daño subletal causados por alta presión

Jordi Saldo Periago

CERPTA-UAB
www.cerpta.com




TECNIO
Be tech. Be competitive



Effects of high pressure on macromolecules

- Most of pressure effects on macromolecules are transient
 - Changes on quaternary and tertiary protein conformation
 - Depolymerisation
 - Lipid crystallisation


Examples of the effects of high hydrostatic pressure on cells and cellular components



A: lipids in membranes
 C: protein



B: multimeric protein assemblages
 E: protein translation by ribosomes

Membrane




http://en.wikipedia.org/wiki/Image:Cell_membrane_detailed_diagram.svg

Protein synthesis system



Cell division







Microbial sub-populations after pressure treatment

- Healthy
- Stressed
- Injured
- Dead




Techniques to distinguish between states


- Dyes
 - Flow cytometry



- Differential growth




- Lag phase



Matrix

Tris Buffer





Dades



☒ Dead cells

☒ Injured cells

☒ Stressed cells

☒ Healthy cells


Code

Storage Days





Matrix
 Skim Milk



Dades


- Dead cells
- Injured cells
- Stressed cells
- Healthy cells

Code
 Storage Days

Matrix

Orange Juice





Dades



☒ Dead cells
☒ Injured cells
☒ Stressed cells
☒ Healthy cells

Code

Storage Days




SISTEMAS ACUOSOS II. DISOLUCIONES

Pedro D. Sanz

Lab. Alta Presión Hidrostática en Alimentos
Departamento de Procesos
Instituto de Ciencia y Tecnología de los Alimentos y Nutrición (ICTAN)
Consejo Superior de Investigaciones Científicas
C/ José Antonio Nováis, 10
Ciudad Universitaria
E-28040 Madrid



Miraflores de la Sierra, 24 de Enero de 2010

Materias primas



ALTA
SEGURIDAD MICROBIOLÓGICA



PROCESADO DE ALIMENTOS

MÍNIMA ALTERACIÓN
CARACTERÍSTICAS
ORGANOLÉPTICAS Y NUTRICIONALES

PROLONGACIÓN
VIDA ÚTIL


Productos elaborados




Secado Cocción
Congelación

Esterilización
Pasterización

Pascalización




Homo Sapiens
- 25 000



Appert
1800




Bridgman
1930



Evolución de las tecnologías de
conservación de alimentos




0.1MPa, 40°C, 20min




Almidón
de trigo

400MPa, 40°C, 20min



■ TIPOS DE PROCESOS



➤ **Conservación en estado no congelado** a temperaturas menores de 0°C

➤ **Congelación** con alta presión

➤ **Descongelación** con alta presión

ALTA PRESIÓN HIDROSTÁTICA:

Tecnología **emergente**


Prolongación vida útil de los alimentos

Inactivación microbiana y/o enzimática

Bajo consumo energético

Respeto al **medio ambiente**

Number of HPP installations in the world



SISTEMAS ACUOSOS II. DISOLUCIONES


I. WATER AND LIFE-RELATED SYSTEMS

**b. High pressure effects on aqueous solutions:
salts, supramolecular aggregates and proteins**

Now!

ALTAS PRESIONES

[Rasanayagam y col., 2003]



Aumento de temperatura durante la compresión





HIGH PRESSURE PROCESSING

Pascal's Law

Instantaneous


$$\frac{dT}{dP} = \frac{\alpha \cdot T_k \cdot V}{C_p}$$

slow



Otero, L.; Molina-García, A. D. & Sanz, P. D.
"Thermal effect in foods during quasi-adiabatic pressure treatments" *Innovative Food Science & Emerging Technologies*, 1 (2), 119-126. 2000



Expansion of 50 MPa at initial conditions of 30°C and 50




Tecnologías combinadas

PROCESADO TÉRMICO ASISTIDO POR PRESIÓN

Uniformidad térmica ESENCIAL



Inactivación de esporas de *Clostridium sporogenes* (PA 3679) en leche tratada a 900 MPa/ 100°C



Objetivos conseguidos

SOLID SAMPLE

$$\rho C_p \frac{\partial T}{\partial t} - \nabla \cdot (k \nabla T) = Q_p$$

$$\frac{dT}{dP} = \frac{\alpha \cdot T_k \cdot V}{C_p}$$

PRESSURIZATION FLUID

$$\rho \frac{\partial \mathbf{u}}{\partial t} - \nabla \cdot \eta (\nabla \mathbf{u} + (\nabla \mathbf{u})^T) + \rho (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla P = \mathbf{g} \rho$$

$$\nabla \cdot (\rho \mathbf{u}) = 0$$

$$\rho C_p \frac{\partial T}{\partial t} + \nabla \cdot (-k \nabla T + \rho C_p \mathbf{u} T) = Q_p$$

Control
150 MPa
200 MPa
250 MPa
300 MPa
350 MPa
400 MPa
450 MPa
500 MPa
550 MPa
600 MPa
Control



Estructura
primaria



Secuencia
de aminoácidos

Estructura
secundaria




Plegamiento local
determinado por
interacciones de
corto alcance


Estructura
terciaria



Plegamiento adicional
determinado por fuerzas
de largo alcance



Un polipéptido con tres
conformaciones de
distinto tiempo de vida



Colaboración
con UCM-
SIMPOL

b. High pressure effects on aqueous solutions of supramolecular aggregates and proteins

Data on the physical properties of food and food components are still scarce at high pressure. These data are needed to develop and improve food high pressure processes.


Pressure dependence of density, isothermal compressibility, thermal expansivity, specific heat and thermal conductivity are required for numerical simulation of heat and mass transfers and of the coupled microorganism inactivation during high pressure treatment. In this way, treatment uniformity can be determined.

Pressure effects on protein conformation and on aggregation phenomena can be assessed by volumetric and ultrasounds techniques *in situ*. The apparent properties of food components in aqueous solution provide information about their behaviour with pressure in foods. Speed of sound also allows for the determination of critical micellar concentration under pressure.



Specific volume and isothermal compressibility were determined for a wide variety of liquid foods and pressure transmitting fluids: pure water, tomato paste, orange juice, whole and skim milk, sodium caseinate aqueous solutions, vegetable oils, silicone oil, castor oil, propylene glycol, ethylene glycol and ethanol.

Ultrasonic measurements were performed in tomato paste and sodium laurate aqueous solutions.




References

- Otero et al. 2010. Modeling thermophysical properties of food under high pressure. *Crit. Rev. Food Sci.*, 50(4), 344-368.
- Guignon et al. 2010. Volumetric properties of pressure-transmitting fluids up to 350 MPa: water, ethanol, ethylene glycol, propylene glycol, castor oil, silicone oil and some of their binary mixture. *J. Chem. Eng. Data*, 55, 3017-3023.
- Guignon et al. 2010. Specific volume of liquid water from (253.15 to 323.15) K and pressures up to 350 MPa by volumetric measurements. *J. Chem. Eng. Data*, 55, 3338-3345.
- Aparicio et al. 2011. Thermal expansion coefficient and heat capacity from sound velocity measurements in tomato paste from 0.1 MPa up to 350 MPa as a function of temperature. *J. Food Eng.*, submitted.

Evaluación de incertidumbres

$$v(p) = v_{ref} + \Delta v(p)$$



$$u_c(T) = \sqrt{u^2(T_s) + u^2(T_{cal}) + u^2(T_{rep})}$$

$$u_c(p) = \sqrt{u^2(p_s) + u^2(p_{cal}) + u^2(p_{rep})}$$

$$u_c(v) = \sqrt{u^2(v_{ref}) + u_c^2(\Delta v) + u^2(v_{uT}) + u^2(v_{up})}$$

Figure 1. Volumetric device: 1, high pressure hydraulic pump; 2, high pressure vessel; 3, thermostatic bath; 4, LVDT; 5, solid piston; 6, sample.

$$u_c(\Delta v) = \sqrt{\left(\frac{\partial(\Delta v)}{\partial k}\right)^2 \cdot u^2(k) + \left(\frac{\partial(\Delta v)}{\partial U_0}\right)^2 \cdot u^2(U_0) + \left(\frac{\partial(\Delta v)}{\partial U(p)}\right)^2 \cdot u^2(U(p)) + \left(\frac{\partial(\Delta v)}{\partial r}\right)^2 \cdot u^2(r) + \left(\frac{\partial(\Delta v)}{\partial m}\right)^2 \cdot u^2(m)}$$

Guignon, Bérengère; Aparicio, Cristina; Sanz, Pedro D.

“Specific Volume of Liquid Water from (253 to 323) K and Pressures up to 350 MPa by Volumetric Measurements”.

Journal of Chemical & Engineering Data, 55, 3338–3345. 2010


High Pressure in Food Lab
ICTAN-CSIC

Influencia de la presión en la simulación del plegamiento de proteínas



Ramiro Perezzan Rodriguez



D




N








D



N



ESTRUCTURA
PDB




ESTRUCTURA
 C_{α}


ESTUDIO
TERMODINAMICO DEL
PROCESO DE
PLEGAMIENTO
MEDIANTE
SIMULACION POR EL
METODO DE
MONTECARLO




D




N






ESTRUCTURA
PDB




ESTRUCTURA
 C_{α}



ESTUDIO
TERMODINAMICO DEL
PROCESO DE
PLEGAMIENTO
MEDIANTE
SIMULACION POR EL
METODO DE
MONTECARLO



ESTRUCTURA
PDB




ESTRUCTURA
 C_{α}



ESTUDIO
TERMODINAMICO DEL
PROCESO DE
PLEGAMIENTO
MEDIANTE
SIMULACION POR EL
METODO DE
MONTECARLO

¿Cómo afecta la presión al plegamiento de proteínas?



¿Cómo afecta la presión al plegamiento de proteínas?



¿Cómo afecta la presión al plegamiento de proteínas?



Boonchai B. Boonyaratanakornkit, Chan Beum Park, Douglas S. Clark *Biochimica et Biophysica Acta* 1595 (2002) 235-249



MovieMaker: a web server for rapid rendering of protein motions and interactions


Rajarshi Maiti^{1,2}, Gary H. Van Domselaar^{1,2} and David S. Wishart^{1,2,*}

¹Department of Computing Science and ²Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E8

high pressure:

147 l.r. cts.

$R_g = 11.28 \text{ \AA}$




1V80-1 (ubiquitin at low pressure: 30 bar)


low pressure:

156 l.r. cts.


$R_g = 11.34 \text{ \AA}$




GB1 domain of protein G




GB1 domain of protein G




Ubiquitin



Ubiquitin





POTENCIALES DE CAMPO MEDIO PARA INTERACCIÓN DE RESIDUOS HIDROFÓBICOS BAJO PRESIÓN



Tuhin Ghosh, Angel E. García and Shekhard Garde

J. Am. Chem. Soc. **2001**, 123, 10997-11003


POTENCIALES DE CAMPO MEDIO PARA INTERACCIÓN DE RESIDUOS HIDROFÓBICOS BAJO PRESIÓN




Tuhin Ghosh, Angel E. García and Shekhard Garde
J. Am. Chem. Soc. **2001**, 123, 10997-11003


Zhirong Liu and Hue Sun Chan
J. Mol. Bio. **2005**, 349, 872-889

$$\epsilon_{\text{go}} - 1.0; \epsilon_{\text{db}} 0.35; \epsilon_{\text{ssm}} - 0.20;$$



$$\epsilon_{\text{go}} - 0.9; \epsilon_{\text{db}} 0.15; \epsilon_{\text{ssm}} - 0.35;$$






$\epsilon_{go}-1.0; \epsilon_{db}0.0; \epsilon_{ssm}-0.35$

$\epsilon_{go}-1.0; \epsilon_{db}0.5; \epsilon_{ssm}-0.35$

$\epsilon_{go}-0.9; \epsilon_{db}0.0; \epsilon_{ssm}-0.35$

$\epsilon_{go}-0.9; \epsilon_{db}0.5; \epsilon_{ssm}-0.35$




$\epsilon_{go}-1.0; \epsilon_{db}0.5; \epsilon_{ssm}0.0$

$\epsilon_{go}-1.0; \epsilon_{db}0.5; \epsilon_{ssm}-0.5$

$\epsilon_{go}-0.9; \epsilon_{db}0.5; \epsilon_{ssm}0.0$

$\epsilon_{go}-0.9; \epsilon_{db}0.5; \epsilon_{ssm}-0.5$








$\epsilon_{go}-1.0; \epsilon_{db}0.0; \epsilon_{ssm}0.0$

$\epsilon_{go}-1.0; \epsilon_{db}0.5; \epsilon_{ssm}0.0$

$\epsilon_{go}-1.0; \epsilon_{db}0.0; \epsilon_{ssm}-0.5$

$\epsilon_{go}-1.0; \epsilon_{db}0.5; \epsilon_{ssm}-0.5$







$\epsilon_{go}-0.84; \epsilon_{db}0.35; \epsilon_{ssm}-0.03$

$\epsilon_{go}-0.90; \epsilon_{db}0.35; \epsilon_{ssm}-0.15$

$\epsilon_{go}-0.96; \epsilon_{db}0.35; \epsilon_{ssm}-0.26$

$\epsilon_{go}-1.01; \epsilon_{db}0.35; \epsilon_{ssm}-0.37$

$\epsilon_{go}-1.07; \epsilon_{db}0.35; \epsilon_{ssm}-0.50$




$\epsilon_{go}-0.84; \epsilon_{db}0.35; \epsilon_{ssm}-0.03$



$\epsilon_{go}-0.90; \epsilon_{db}0.35; \epsilon_{ssm}-0.15$

$\epsilon_{go}-0.96; \epsilon_{db}0.35; \epsilon_{ssm}-0.26$

$\epsilon_{go}-1.01; \epsilon_{db}0.35; \epsilon_{ssm}-0.37$

$\epsilon_{go}-1.07; \epsilon_{db}0.35; \epsilon_{ssm}-0.50$







$\epsilon_{go}-0.84; \epsilon_{db}0.35; \epsilon_{ssm}-0.03$

$\epsilon_{go}-0.90; \epsilon_{db}0.35; \epsilon_{ssm}-0.15$

$\epsilon_{go}-0.96; \epsilon_{db}0.35; \epsilon_{ssm}-0.26$

$\epsilon_{go}-1.01; \epsilon_{db}0.35; \epsilon_{ssm}-0.37$

$\epsilon_{go}-1.07; \epsilon_{db}0.35; \epsilon_{ssm}-0.50$



GRUPO UCM-SIMPOL

Prof. Antonio Rey Gayo

Prof. Ana María Rubio Caparrós

Dra. María Larriva Hormigos

Lcda. Marta Enciso Carrasco

Lcdo. Ramiro Perezzan Rodríguez



QUIM  PRES






Nanoencapsulació por Ultra Alta Presión Homogenización (UHPH)


Roger Escriu Justo

Formación “natural” nanoestructuras

Leche cruda




300 MPa (30°C)




Micelas caseina
Globulos grasos

Formación “natural” nanoestructuras

Leche cruda






300 MPa (30°C)



Micelas caseina

Globulos grasos


¿Qué es nanotecnología?



Es un conjunto de técnicas que se utilizan para manipular la materia a la escala de **átomos y moléculas**.

Del griego νᾶνος
(enano), τέχνη
(técnica) y λόγος
(ciencia)

A escala tan pequeña, la materia se comporta de modo diferente.




Richard P. Feynman

(precursor de la nanotecnología):

“¿Qué pasaría si nosotros pudiéramos arreglar los átomos uno por uno de la manera en que nosotros los queremos?”

(1960)




© Copyright California Institute of Technology. All rights reserved.
Commercial use or modification of this material is prohibited.

La **NANOTECNOLOGÍA** constituye un ensamblaje interdisciplinar de **varios** campos de las ciencias naturales que están altamente especializados.

FÍSICA: En la construcción de los microscopios usados para investigar tales fenómenos y por su conocimiento de las leyes de la mecánica cuántica.


QUÍMICA: Conocimiento de la estructura del material deseado y las configuraciones de ciertos átomos




Ultra Alta Presión de Homogeneización (UHPH)

Primeros Homogeneizadores

1900



1902. Gaulin



Objetivos homogeneizadores convencionales

- Estabilización de emulsiones
- Mejora del sabor
- Mejora de la textura
- Leche, nata y helados
- Presiones hasta 50 MPa

1980 nace la alta presión homogeneización (HPH)


- Presiones hasta 150 MPa
- Producir emulsiones más estables
- Mejorar la textura de los yogures
- Reducir algo la carga microbiana

Alta Presión Homogeneización (HPH)




Hasta 1500 bares
(150 MPa)


1990. Primeros prototipos de ultra alta presión homogenización (UHPH)




2003. Primeras válvulas y prototipos de UHPH con doble intensificador capaces de soportar 400 MPa



Equipo Ultra Alta Presión Homogeneización. 8 L/h




Válvula Homogeneización



Incremento de temperatura:

El incremento lineal de temperatura se produce al incrementar la presión de homogenización

- Calentamiento adiabático debido al incremento en la presión
- Alta velocidad, turbulencia y fuerzas de corte al pasar y salir de la válvula




$19.5 \pm 3 \text{ }^{\circ}\text{C}$ por 100 MPa

(Pereda et. al, 2006; Thiebaud et al., 2003; Hayes et al., 2003; Flourey et al., 2000)





Ligero efecto térmico: Tiempo de residencia 10^{-4} s *(Flourey et al., 2002)*



F. Donst et al. / Chemical Engineering Science 64 (2009) 520–532

Equipo Ultra Alta Presión Homogeneización. 120 L/h




Aplicaciones en Alimentos


- Pasteurización y esterilización de alimentos fluidos
- Control de modificaciones sobre las proteínas
- Mejores texturas e incremento de la calidad
- Desarrollo de nuevos productos
- Ventajas:
 - Sistema continuo. Posibilidad de envasado aséptico
 - Economía de energía
 - Sistema higiénico de fácil limpieza y desinfección

1. Presiones de trabajo entre 100 hasta 400 MPa.
 - Gracias a nuevos desarrollos en la ciencia de los materiales
 - Diversas empresas constructoras y nuevos equipos: Microfluidizer®, Emulsiflex®, Stansted Fluid Power
2. Cavitación, turbulencia y fuerzas de cizalla.
3. Aplicaciones actuales:
 - **Elaboración de nanodispersiones y nanocápsulas**
 - Extracción de enzimas de las células microbianas
4. Potenciales aplicaciones:
 - Tratamiento de alimentos líquidos para incrementar su seguridad y sus propiedades funcionales.


Estructura esquemática de los diferentes “nanoportadores”




Liposome




Nanosphere



Nanocapsule






Dendrimer



Micelle





Inflamación

División
celular y
crecimiento

Apoptosis

Reparación
DNA

Regulación
Hormonal

Metabolismo



Balances
energéticos

Componentes Bioactivos en Alimentos

- Vitaminas
- Minerales
- Lípidos funcionales
- Bacterias Probióticas
- Amino acidos
- Proteinas
- ...

La encapsulación protege ingredientes alimentarios sensibles al calor, humedad y pH.

¿Cómo se hacen las nanocápsulas?



Nanotecnología y alimentación



NANOPARTÍCULAS comestibles para:

Crear bebidas inteligentes
(con sabores y colores
específicos).

“Mejorar” ciertos
alimentos
(nanoencapsulando
antioxidantes para
aumentar la absorción de
alimentos específicos).

Hacer fármacos más
efectivos.



“Tip-Top Up” – Pan con Omega 3

- Nanocápsulas con Omega 3
- Nanocápsulas que se rompen únicamente en el estómago




Tip-Top bakery - Australia

Otras aplicaciones de la Nanotecnología...




**Pinturas protectoras
contra la corrosión**



**Raquetas de tenis más
fuertes y ligeras**



**Tejidos resistentes a
las manchas**



[Add to Cart](#)

**Pelotas de tenis
de largo duración**



**Protectores solares y
cosméticos**



Simposio
MALTA-Consolider

23-27 Enero 2011, Miraflores de la Sierra



www.malta-consolider.com

Muchas gracias!!

Efectos inducidos por la presión sobre agregados en medio acuoso:



Aplicación de Técnicas Volumétricas al Estudio de la Asociación de Caseínas



Bérengère Guignon

Instituto de Ciencia y Tecnología de los Alimentos y Nutrición



Las caseínas





Las altas presiones permiten modificar las propiedades tecnológicas de las caseínas

0,1 MPa

200 MPa


400 MPa



Dispersión de luz



Microscopía



Objetivo:

evaluar la aplicación de técnicas volumétricas para obtener información sobre el estado de agregación de las caseínas

Caracterización de propiedades
volumétricas a P_{atm}



Estrategia

Medidas macro

v_ϕ $k_{s\phi}$



Medidas micro

Datos publicados



Caracterización a alta presión

Medidas volumétricas

Patm

Alta Presión

$$1/\rho = v$$

Densidad ρ $\leftrightarrow$ Volumen específico **v**

+ Velocidad del sonido u_s

$$v/u_s^2 = k_s$$

$\rightarrow$ Compresibilidad adiabática **k_s**

Volumen específico aparente

$$v_{\phi} = \frac{1}{w} (v - v_0) + v_0$$

Disolución de caseínas agua

$$v_{\phi} = \underbrace{v_{\text{átomos}}}_{\text{white}} + \underbrace{v_{\text{cavidad}}}_{\text{yellow}} + \underbrace{v_{\text{hidratación}}}_{\text{blue}} + \underbrace{v_{\text{térmico}}}_{\text{red}} + v_{\text{ideal}}$$



Compresibilidad adiabática aparente


$$k_{s\phi} = \frac{1}{w} \left(\frac{v^2}{u_s^2} - \frac{v_0^2}{u_{s0}^2} \right) + \frac{v_0^2}{u_{s0}^2}$$



Disolución de caseínas   agua

$$k_{s\phi} = k_{s\text{ cavidad}} + k_{s\text{ hidratación}} + k_{s\text{ térmico}}$$



Resultados

A Presión Atmosférica






v_ϕ a presión atmosférica



Δv_ϕ a alta presión



Conclusión

A presión atmosférica:

- el proceso de asociación de las caseínas no es cooperativo
- las técnicas volumétricas son válidas para el estudio del estado de agregación de las caseínas.

A alta presión, se han identificado las siguientes dificultades para el desarrollo del método volumétrico:

- repetibilidad de p y T de un ensayo a otro (solvente, velocidad del sonido)
- incertidumbre de la presión

Efectos inducidos por la presión sobre agregados en medio acuoso. Estudio del dodecanoato de sodio mediante medidas de velocidad del sonido



Eduardo Hidalgo Baltasar



Introducción

Procesado de alimentos a alta presión



Sabor, olor, color

Textura



Microbiología




Medio Ambiente

Economía





Introducción

Monómeros anfifílicos




Micelas

Las micelas como modelos de sistemas biológicos



Introducción

Concentración Micelar Crítica (cmc)




Cambio de Pendiente

cmc

Introducción

- ¿Por qué las membranas celulares se vuelven permeables?
- ¿Qué ley rige el efecto de la presión sobre las interacciones débiles?
- ¿Por qué las micelas de caseína se disocian a 150-250 MPa?



Introducción



- UCM
 - Dpto. Química Física I
 - Espectroscopía Raman
 - Altas presiones: hasta 5 GPa






- CSIC-ICTAN
 - Dpto. de Procesos
 - Propiedades volumétricas y acústicas bajo presión
 - 45 mL - 700 MPa

Efecto de la presión en sistemas micelares

Objetivos

- Efecto de la presión en diversas propiedades físico-químicas del sistema micelar agua-dodecanoato de sodio




Experimentos

Densidad

Ultrasonidos

Espectroscopía
Raman



Experimentos




Ultrasonidos

A Presión atmosférica

Alta Presión hasta 700 MPa

Dispositivos experimentales


- Transmisión con piezoeléctricos de 2 MHz






$$u_s = \frac{2d}{\Delta t}$$




Tratamiento de la señal





Transformada
de Fourier



Tratamiento de la señal




Transformada
de Fourier





Leer tiempos con precisión

Determinación del tiempo de vuelo




$$\Delta t = t_2 - t_1$$

$$u_s = \frac{2d}{\Delta t}$$





Equipo de Alta Presión





Resultados experimentales:

Tiempo de vuelo





Resultados experimentales:

Velocidad del sonido a presión atmosférica




$$u_s = \frac{2d}{\Delta t}$$

Resultados experimentales: Velocidad del sonido a altas presiones



Conclusiones

- Presión atmosférica y alta presión



Validez de la técnica de ultrasonidos

- Presión atmosférica

Densidad



Volumen molar aparente

Densidad + Ultrasonidos



Compresibilidad Adiabática

Gracias por la atención



MALTA Reunión, Miraflores, España

El enlace químico en moléculas heterodiatómicas bajo la perspectiva de la teoría de Átomos Cuánticos Interaccionantes (IQA)

Aurora Costales, M. A. Blanco, E. Francisco y
A. Martín Pendás



Universidad de Oviedo
Dpto. Química Física y Analítica

Organización de la charla

1. Partición energética QTAIM
2. La teoría IQA
3. Energía de enlace en moléculas heterodiatómicas
4. Conclusiones

Organización de la charla

1. Partición energética QTAIM
2. La teoría IQA
3. Energía de enlace en moléculas heterodiatómicas
4. Conclusiones

Organización de la charla

1. Partición energética QTAIM
2. La teoría IQA
3. Energía de enlace en moléculas heterodiatómicas
4. Conclusiones

Organización de la charla

1. Partición energética QTAIM
2. La teoría IQA
3. Energía de enlace en moléculas heterodiatómicas
4. Conclusiones

Partición energética QTAIM

- Partición del espacio en cuencas atómicas:

$$R^3 \rightarrow \bigcup_A \Omega_A \Rightarrow 1 = \sum_A \omega_A, \quad \omega_A(\vec{r}) = \begin{cases} 1 & \vec{r} \in \Omega_A \\ 0 & \vec{r} \notin \Omega_A \end{cases}$$

Partición energética QTAIM

- Partición del espacio en cuencas atómicas:

$$R^3 \rightarrow \bigcup_A \Omega_A \Rightarrow 1 = \sum_A \omega_A, \quad \omega_A(\vec{r}) = \begin{cases} 1 & \vec{r} \in \Omega_A \\ 0 & \vec{r} \notin \Omega_A \end{cases}$$

- Partición de las matrices densidad:

$$\rho(\vec{r}) = \sum_A \rho_A(\vec{r}) \text{ con } \rho_A(\vec{r}) = \rho(\vec{r})\omega_A(\vec{r}).$$

Particiones similares para $\rho_1(1; 1')$ y $\rho_2(1, 2)$.

Se puede extender: Cuencas ELF, átomos difusos, ...

Partición energética QTAIM

- Partición del espacio en cuencas atómicas:

$$R^3 \rightarrow \bigcup_A \Omega_A \Rightarrow 1 = \sum_A \omega_A, \quad \omega_A(\vec{r}) = \begin{cases} 1 & \vec{r} \in \Omega_A \\ 0 & \vec{r} \notin \Omega_A \end{cases}$$

- Partición de las matrices densidad:

$$\rho(\vec{r}) = \sum_A \rho_A(\vec{r}) \text{ con } \rho_A(\vec{r}) = \rho(\vec{r})\omega_A(\vec{r}).$$

Particiones similares para $\rho_1(1; 1')$ y $\rho_2(1, 2)$.

Se puede extender: Cuencas ELF, átomos difusos, ...

- Así se induce una partición de la energía total

$$E = E_1 + E_2 + \frac{1}{2} \sum_{A,B \neq A} V_{nn}^{AB} \text{ con}$$

$$E_1 = \sum_A T^A + \sum_{A,B} V_{en}^{AB}$$

$$E_2 = \sum_A V_{ee}^{AA} + \frac{1}{2} \sum_{A,B \neq A} V_{ee}^{AB}$$

Teoría IQA: términos de 1 & 2-átomos

Términos energéticos: T^A , V_{nn}^{AB} , $V_{en}^{AB} = V_{ne}^{BA}$, V_{ee}^{AB}

$A = B$ (1 átomo), $A \neq B$ (2 átomos)

Teoría IQA: términos de 1 & 2-átomos

Términos energéticos: T^A , V_{nn}^{AB} , $V_{en}^{AB} = V_{ne}^{BA}$, V_{ee}^{AB}

$A = B$ (1 átomo), $A \neq B$ (2 átomos)

- $E_{\text{net}}^A = T^A + V_{en}^{AA} + V_{ee}^{AA}$,
equivalente a las energías de los átomos libres

Teoría IQA: términos de 1 & 2-átomos

Términos energéticos: T^A , V_{nn}^{AB} , $V_{en}^{AB} = V_{ne}^{BA}$, V_{ee}^{AB}

$A = B$ (1 átomo), $A \neq B$ (2 átomos)

- $E_{\text{net}}^A = T^A + V_{en}^{AA} + V_{ee}^{AA}$,
equivalente a las energías de los átomos libres
- $E_{\text{int}}^{AB} = V_{nn}^{AB} + V_{en}^{AB} + V_{ne}^{AB} + V_{ee}^{AB}$,
incluye todas las interacciones interatómicas
interpartículas

Teoría IQA: términos de 1 & 2-átomos

Términos energéticos: T^A , V_{nn}^{AB} , $V_{en}^{AB} = V_{ne}^{BA}$, V_{ee}^{AB}

$A = B$ (1 átomo), $A \neq B$ (2 átomos)

- $E_{\text{net}}^A = T^A + V_{en}^{AA} + V_{ee}^{AA}$,
equivalente a las energías de los átomos libres
- $E_{\text{int}}^{AB} = V_{nn}^{AB} + V_{en}^{AB} + V_{ne}^{AB} + V_{ee}^{AB}$,
incluye todas las interacciones interatómicas interpartículas
- $E = \sum_A E_{\text{net}}^A + \frac{1}{2} \sum_{A,B \neq A} E_{\text{int}}^{AB}$,
incluye las energías de los **átomos cuánticos** más una suma de sus interacciones de pareja

Energías de enlace y de deformación

- E_{bind} se define respecto a una referencia

$$E_{\text{bind}} = E - \sum_A E_{\text{vac}}^A$$

Energías de enlace y de deformación

- E_{bind} se define respecto a una referencia

$$E_{\text{bind}} = E - \sum_A E_{\text{vac}}^A = \sum_A E_{\text{def}}^A + \frac{1}{2} \sum_{A,B \neq A} E_{\text{int}}^{AB}$$

- Definimos la E_{def} como la diferencia entre las energías atómicas **en la molécula** y **en el vacío**

$$E_{\text{def}}^A = E_{\text{net}}^A - E_{\text{vac}}^A$$

Energías de enlace y de deformación

- E_{bind} se define respecto a una referencia

$$E_{\text{bind}} = E - \sum_A E_{\text{vac}}^A = \sum_A E_{\text{def}}^A + \frac{1}{2} \sum_{A,B \neq A} E_{\text{int}}^{AB}$$
- Definimos la E_{def} como la diferencia entre las energías atómicas **en la molécula** y **en el vacío**

$$E_{\text{def}}^A = E_{\text{net}}^A - E_{\text{vac}}^A \equiv E_{\text{def}}^A(\text{CT}) + E_{\text{def}}^A(\text{CR})$$
 - Transferencia de carga, $E_{\text{def}}^A(\text{CT}) > 0$ para $Q^A > 0$ (catión), negativa para el anión
 - Redistribución de carga: $E_{\text{def}}^A(\text{CR}) > 0$
- En general, $E_{\text{def}}^A > 0 \Rightarrow \sum E_{\text{def}}^A > 0 \Rightarrow \sum E_{\text{int}}^{AB} < 0$ y para que exista enlace:
compensación entre deformaciones pequeñas e interacciones estabilizantes grandes

Interacciones: Coulomb & exch-corr

- Definiendo $\rho_2^C(1, 2) = \rho(1)\rho(2)$ y $\rho_2^{xc} = \rho_2 - \rho_2^C$:

$$V_{ee}^{AB} = V_C^{AB} + V_{xc}^{AB}$$

- Definiendo $V_{cl}^{AB} = V_C^{AB} + V_{en}^{AB} + V_{ne}^{AB} + V_{nn}^{AB}$:




$$E_{int}^{AB} = V_{cl}^{AB} + V_{xc}^{AB}$$

- Para densidades bien-separadas, $V_{cl}^{AB} \simeq$ multipolar, $E_Q^{AB} = Q^A Q^B / R_{AB} \Rightarrow$ ionicidad
- $V_{xc}^{AB} = \int_A \int_B \rho_2^{xc} r_{12}^{-1}$ relacionada con la covalencia. El orden de enlace se mide con $\delta_{AB} = 2 \left| \int_A \int_B \rho_2^{xc} \right|$, que determina el número de pares de electrones compartidos

Detalles computacionales




- Heterodiatómicas: series isoelectrónicas, $12e$ (LiF, BeO, BN, C_2), $14e$ (BF, CO, N_2), y $20e$ (NaF, MgO, AlN, SiC)
- TZV(d) cálculos CAS: $[8, 8]$ (iso12, 20), y $[10, 10]$ (iso14)
- 1-2 % error en R_e , 5-10 % en D_e con respecto a valores experimentales
- Integraciones IQA 2- e : $600(r) \times 3890$ (Lebedev) + β -esferas
- Errores en $Q < 1 m_e$ y $0.1 mE_h$ en las energías
- Unidades: Q en e , E en E_h

Ionicidad y cargas atómicas



Q^A






Ionicidad y cargas atómicas







Q^A Q^A/Q_{nom}^A

La ionicidad se define, en general, comparando con los hipotéticos iones nominales




Componentes de la Energía de enlace



Componentes de la Energía de enlace



Componentes de Deformación ($Q^A > 0$)






$$E_{\text{def}}^A = (\text{CT}) + (\text{CR})$$

$$(\text{CT}) = \sum \text{IP}_i \times q_i$$

$Q^B < 0$: $(\text{CT}) = \sum \text{AE}_i \times q_i$
 difícil calcular AE; son muy polarizables $\Rightarrow$ CR alta

Componentes de Interacción






$$E_{\text{int}}^{AB} = V_{\text{cl}}^{AB} + V_{\text{xc}}^{AB}$$

$$V_{\text{cl}}^{AA} > 0 \text{ in } A_2$$

La covalencia está principalmente unida a V_{xc}^{AB}




Interacción Clásica



V_{cl}^{AB} (white bar) E_Q^{AB} (red bar)
 $V_{cl}^{AB} - E_Q^{AB}$ (green bar) (en general < 0 ; LiF, NaF $\gtrsim 0$)

E_Q^{AB} / V_{cl}^{AB} mide el grado de ionicidad

Energía de Cambio-correlación



■ $-V_{xc}^{AB}$ ($V_{xc}^{AB} < 0$ siempre)

■ δ_{AB} (pares e^- compartidos)

La covalencia está principalmente unida a V_{xc}^{AB} o δ_{AB}

Conclusiones

- IQA permite una descomposición energética de E_{bind} con sentido físico
- La energía de enlace es un balance entre un valor positivo de la deformación (preparación) y uno más negativo de la energía de interacción
- Las huellas de la ionicidad: gran $E_{\text{def}}^A(\text{CT})$ y $E_Q^{AB} / V_{\text{cl}}^{AB}$
- El orden de enlace (δ_{AB}) y V_{xc}^{AB} correlacionan con la covalencia
- Ambos caracteres pueden estar presentes en el mismo enlace en diferentes grados, pero no son complementarios! (Ej., CO)

Further equations

- $\rho_1 = \sum_A \rho_1^A, \quad \rho_2 = \sum_{A,B} \rho_2^{AB}$
- $T^A = \int_\infty \hat{T} \rho_1^A = \int_A \hat{T} \rho_1$
- $V_{en}^{AB} = V_{ne}^{BA} = \int_A -\frac{Z_B}{r_{1B}} \rho$
- $V_{nn}^{AB} = Z_A Z_B / R_{AB} \quad (A \neq B)$
- $V_{ee}^{AA} = \frac{1}{2} \int_A \int_A \rho_2 r_{12}^{-1} \quad (A = B)$
- $V_{ee}^{AB} = \int_A \int_B \rho_2 r_{12}^{-1} \quad (A \neq B)$
- $\rho_2^C(1, 2) = \rho(1)\rho(2)$
- $\rho_2^X(1, 2) = -\rho_1(2; 1)\rho_1(1; 2)$
- $\rho_2 = \rho_2^C + \rho_2^{xc} = \rho_2^C + \rho_2^X + \rho_2^{\text{corr}}$



UNIVERSITAT
JAUME•I

INCORPORANDO LA PRESIÓN HIDROSTÁTICA EN SIMULACIONES DE ESTRUCTURA Y REACTIVIDAD QUÍMICA EN SISTEMAS CONDENSADOS. MÉTODOS QM/MM+DM

José Javier Ruiz Pernía, Juan Andrés Bort.

Reunión proyecto MALTA, Miraflores, Madrid

24/Enero/2010



INTRODUCCIÓN

MÉTODOS QM/MM + DM

- La mayoría de los sistemas y reacciones químicas tienen lugar en presencia de un entorno (en fase condensada).
 - Procesos en disolución.
 - Reacciones enzimáticas, en sólidos, etc
- La fase condensada (entorno) juega un papel fundamental en la termodinámica y en la cinética química

INTERÉS

¿Cómo influye ese medio condensado en la estructura y reactividad química?

Métodos QM
(Cuánticos)

Métodos MM
(Clásicos)

PROBLEMA

Buena precisión

Sistemas pequeños

Nº átomos < 50

Los medios condensados se caracterizan por un gran

número de átomos. Estudio

complejado. Sistemas grandes


Nº átomos > 100000



UNIVERSITAT
JAUME·I

INTRODUCCIÓN

MÉTODOS QM/MM



QM = Directamente involucrados en la ruptura y formación de enlaces químicos

$$E, \frac{\partial E}{\partial R_i}, \frac{\partial^2 E}{\partial R_i^2}$$

MM = Afectan a la reacción indirectamente

$$E, \frac{\partial E}{\partial R_i}$$

$$\hat{H}_{\text{eff}} = \hat{H}_{\text{QM}} + \hat{H}_{\text{MM}} + \hat{H}_{\text{QM/MM}} + \hat{H}_{\text{BOUND}}$$



UNIVERSITAT
JAUME·I

INTRODUCCIÓN

MÉTODOS QM/MM + DM

#1 *Generación de la Superficie de Energía Potencial (SEP)*

** descripción híbrida cuántica / clásica:*

[CHARMM & DYNAMO]

#2 *Exploración de la SEP*

** búsqueda puntos singulares con métodos flexibles:*

[GRACE / CHARMM & DYNAMO]

#3 *Cálculo de la Energía Libre*

** métodos simulación estadística:*




[DYNAMO]



UNIVERSITAT
JAUME·I


INTRODUCCIÓN

Superficies de Energía Potencial (SEP)



Sistemas Grandes

En Medio Condensado







UNIVERSITAT
JAUME·I

INTRODUCCIÓN

Simulaciones QM/MM + DM



$$\zeta = d(A-B) - d(B-C)$$

$\langle \rho(\xi_{0,i}) \rangle$

Permite el cálculo de
propiedades
termodinámicas y cinéticas

ξ_0



UNIVERSITAT
JAUME I

OBJETIVOS

Simulación del Efecto de la Presión

Incorporar la presión hidrostática en simulaciones de estructura y reactividad química de sistemas químicos en fases condensadas (moléculas en disolución, enzimas, etc..).

Modelización de la propiedades físicas y químicas.

Comparación con datos experimentales


Calcular la influencia de la presión sobre las propiedades termodinámicas y cinéticas (estructura y reactividad química)



UNIVERSITAT
JAUME·I


METODOLOGÍA

¿Presión hidrostática en simulaciones QM/MM + DM?



METODOLOGÍA


¿Presión hidrostática en simulaciones QM/MM + DM?



¿El caso más sencillo: agua?

METODOLOGÍA

Incorporando la presión hidrostática en simulaciones
QM/MM + DM




Simular y
calcular la
densidad
del H₂O en
función de la
 $P_{\text{hidrostática}}$

METODOLOGÍA



UNIVERSITAT
JAUME·I

Presión en QM/MM



1 molécula AM1



**Caja 31.4 Å de lado,
1033 moléculas TIP3P**

**Caja 55.4 Å de lado,
5831 moléculas TIP3P**

Cut-off 10.5, 11.5 y 12.5

Simulación NPT

**Algoritmo de
Berendsen**






UNIVERSITAT
JAUME I

RESULTADOS

Sistema Agua AM1/TIP3P (caja 31.4 Å de lado)






UNIVERSITAT
JAUME·I

RESULTADOS

Sistema Agua AM1/TIP3P (caja 55.4 Å de lado)






UNIVERSITAT
JAUME·I

RESULTADOS

Sistema Agua AM1/TIP3P (caja 31.4 Å de lado)

Función de Distribución Radial






UNIVERSITAT
JAUME·I

RESULTADOS


Sistema Agua AM1/TIP3P (caja 55.4 Å de lado)

Función de Distribución Radial



RESULTADOS

Sistema Agua AM1/TIP3P





UNIVERSITAT
JAUME I

RESULTADOS

DESARROLLO METODÓLOGICO

¿Es posible modificar la ecuación de Berendsen para obtener un comportamiento más realista de los campos de fuerza clásicos?

$$P = 2 * (E_{ke} - \Xi) / 3 * V$$

$$P = X * E_{ke} - Y * \Xi / 3 * V$$

Factor Pre-Energía Cinética


Factor Pre-Virial



UNIVERSITAT
JAUME I

RESULTADOS


Sistema Agua AM1/TIP3P



RESULTADOS

Sistema Agua AM1/TIP3P (caja 31.4 Å de lado)


Radial Distribution Function



RESULTADOS

Sistema Agua AM1/TIP3P (caja 55.4 Å de lado)

Radial Distribution Function





UNIVERSITAT
JAUME•I


REACTIVIDAD QUIMICA



UNIVERSITAT
JAUME·I

RESULTADOS


Paso de Chalcona a Isoflavona



En disolución (agua)

- 30 átomos QM (AM1)
- 17439 átomos TIP3P (5813 moléculas de agua de solvatación)
- Caja de 55.4 Å de lado

En medio enzimático

- 
- 30 átomos QM (AM1)
 - 3231 átomos OPLS-AA (enzima)
 - 47745 átomos TIP3P (15915 moléculas de agua de solvatación)
 - Caja de 79.8 Å de lado




UNIVERSITAT
JAUME·I

RESULTADOS

Reacción Chalcona-Isoflavona en agua (caja 55.4 Å)

Función de Distribución Radial






UNIVERSITAT
JAUME·I

RESULTADOS

Perfiles Energía Libre AM1/MM. Reacción en agua






UNIVERSITAT
JAUME I

RESULTADOS

Potenciales de Fuerza Media AM1/MM. Reacción en agua



Presión(atm)	RC [‡] (Å)	ΔPMF [‡] (Kcal/mol)
1	1.837	26.7
1000	1.839	24.9
5000	1.857	20.3
15000	1.840	19.0
20000	1.863	15.9

CONCLUSIONES

- Se ha desarrollado, implementado y aplicado una metodología QM/MM+DM novedosa para la simular la estructura y reactividad química en fases condensadas, incluyendo la presión hidrostática.
- Se ha obtenido un adecuada representación del campo de fuerzas MM, para reproducir el comportamiento del agua. i.e, la representación de la densidad en función de la presión, utilizando la ecuación de Berendsen modificada.
- Se ha estudiado la reacción de chalcona a isoflavona en agua, obteniéndose los perfiles de energía libre y las estructuras de transición, reactivos y productos en función de la presión hidrostática. En curso, la reacción en medio enzimático.

PERSPECTIVAS DE FUTURO

- Estudiar la influencia de la presión hidrostática en el mecanismo molecular de reacciones orgánicas (S_N2 , S_N1 , E2, Diels-Alder, etc..) en agua
- Desarrollar potenciales, MM, para simular la estructura y reactividad química en otros disolventes.
- Extender la metodología a reacciones químicas en sistemas biológicos (enzimas).
- Aumentar el nivel cuántico de representación de la parte QM (DFT, *ab-initio*, híbrido, etc).
- Estudiar la posible influencia de la presión como coordenada colectiva sobre la coordenada distinguida.



UNIVERSITAT
JAUME•I


AGRADECIMIENTOS

**Juan Andrés Bort,
Armando Beltrán,
Lourdes Gracia Edo
y
Sergio Martí**





Modelización de la alta presión con métodos de función de onda

Reactividad a alta presión del dimetilacetileno



Dimetilacetileno (DMA)










Tabla 4.1: Parámetros geométricos calculados y experimentales [85] del DMA. Las distancias se expresan en Å y los ángulos en grados. Base: 6-31++G(d,p) para B3LYP, MP2, CASSCF y MRMP2 y ondas planas para CASTEP.


Método	$r_{C_1C_3}$	$r_{C_1C_2}$	$r_{C_3H_5}$	$\alpha_{C_3C_1C_2}$	$\alpha_{H_5C_3C_1}$
B3LYP	1.463	1.212	1.096	179.2	111.2
MP2	1.464	1.224	1.090	179.6	110.8
CASSCF(4,4)	1.472	1.200	1.084	179.5	110.7
MRMP2(4,4)	1.471	1.199	1.084	179.3	110.7
Castep	1.434	1.194	1.091	179.8	118.2
Experimental, fase II	1.468	1.201	1.091	178.5	111.4





Tabla 4.2: Frecuencias calculadas y experimentales [26] del DMA (en cm^{-1}). La bases usadas han sido la 6-31++G(d,p) para B3LYP, MP2 y CASSCF y ondas planas para CASTEP/PBE.


Descripción	Modo	Experimental	Castep	B3LYP	MP2	CAS(4,4)
$\equiv \text{C} - \text{C}$ flexión	ν_{16}	390	409	378	206	440
$\text{C} - \text{CH}_3$ tensión	ν_4	723	724	725	730	747
CH_3 rocking	ν_{15}	1048	1046	1055	1062	1103
CH_3 deformación	ν_3	1399	1391	1423	1459	1546
	ν_{14}	1455	1450	1484	1526	1587
$\text{C} \equiv \text{C}$ tensión	ν_2	2290 ¹	2281	2358	2298	2476
CH_3 tensión	ν_1	2946	2954	3031	3117	3193
	ν_{13}	2979	3018	3093	3207	3259





































Modo imaginario a 1554 cm^{-1}
relacionado con la automerización


$$H(q) = -\frac{1}{2\mu} \frac{d^2}{dq^2} + aq^2 + bq^4 + c \exp(-dq^2)$$













Modelo de Drickamer



1. Metodología
MCSCF


2. Cambios
químicos



1. Metodología
monodeterminantal

2. Cambios Físicos

• Correlación P-R



Gracias por su atención

Joint Use of the Catastrophe Theory and the Analysis of the Electron Density (QTAIM, ELF and NCI) in reactivity

J. Andrés¹, S. Berski², and J. Contreras-García³

¹Departamento de Química Física y Analítica, Universitat Jaume I

²Faculty of Chemistry, University of Wrocław

³Department of Chemistry, Duke University


SUMMARY

- Non-covalent interactions have a unique signature and their presence can be revealed solely from the electron density.
- This approach provides a rapid and rich representation of van der Waals interactions, hydrogen-bonds, and steric repulsion.
- It only requires the atomic coordinates as input
- It is applicable to a wide range of systems: small/large, organic/inorganic, solids, etc.
- This tool offers exciting possibilities in design and generation of leads.


Reference: J. Am. Chem. Soc. 132, 6498 (2010).

SMALL MOLECULES


We have recently developed an index for visualizing non covalent interactions



SOLIDS



Diamond




Graphite

BIOMOLECULES

- One of the major areas of application of weak interactions are biomolecules
- HOWEVER, SCF calculations are extremely slow
- Reduced density gradient characteristics are preserved from promolecular to SCF density
- only require input atomic coordinates,
- very fast calculation; applicable to large systems

BIOMOLECULES


Polipeptide: α helix with 15 alanine residues



- Hydrogen bonds stabilize the helix
- Big region of van der Waals interaction inside the helix and between methyl lateral chains one step away


BIOMOLECULES

polypeptide: anti-parallel β -sheet consisting of 17 glycine residues




- HB between C=O and N-H groups in parallel chains
- Van der Waals interactions between CH₂ groups

BIOMOLECULES



Non covalent interactions are also crucial for reactivity

REACTIVITY





- This reaction is a well known method for the utilization of hydrogen-based energy. This is a first step in a simple way for fast ammonia utilization or production of hydrogen storage materials.
- The pathway associated with the intrinsic reaction coordinate has been calculated at the B3LYP/aug-cc-pVTZ level.

REACTIVITY

The mechanism has been thoroughly analyzed using topological analysis of:



- The electron density (Five steps)



REACTIVITY

The mechanism has been thoroughly analyzed using topological analysis of:


- Electron Localization Function (Eight steps)



REACTIVITY

What about non covalent interactions?

AÑADIR AQUÍ LA PELI




Magnetism in solid oxygen under pressure

Stefan Klotz

IMPMC

Université P&M Curie, Paris








A.L. Lavoisier
(1743-1794)


όξύς (oxys) -γενής (-genēs)

« acid former »

Oxygen






AO



AO


1848: Faraday:	Paramagnetic nature of O ₂
1910: Kamerlingh Onnes:	Magn. suscept. of liquid and solid O ₂
1963: Alikhanov:	Magn. structure of solid O ₂ (neutrons, 1 bar)
2004: Goncharenko:	Mag. struct. of δ -O ₂ under pressure (neutrons, 6 GPa)







O₂ at 1 bar

β : short-range ordered





α : antiferromagnetic



α - oxygen







$$b/a = 0.79 \neq 3^{1/2}/2$$




δ - oxygen



Nuclear Structures





Previous high P neutron experiments




Goncharenko, PRL 2004

Infrared absorption (diamond anvil cell)



Gorelli et al., PRB 2008



Sample:
80 mm³

TiZr gasket


cBN

~ 10 GPa for $V = 100 \text{ mm}^3$

~ 30 GPa for $V = 10 \text{ mm}^3$

3.5-320 K

see also talk Domingo Martinez
martes 17:40





D20 high intensity diffractometer
ILL, Grenoble, France

Future:

XTREMED: dedicated high P diffractometer at ILL
(Javier Campo Ruiz miércoles 10:45)

Sample loading



Condensation at 77 K

Gas loader






$P_{\max} = 2000 \text{ bar}$
Compatible with hydrogen



Difficulties

- Texture (imperfect powder) due to the formation of single crystals
- Strong compressibility: $\sim 50\%$ from 1-8 GPa
- Uncontrollable download
- Analysis: anisotropic magnetic form factor







220 K



145 K

100 K



LTC Structure



(Low Temperature Commensurate)




Ferromagnetic coupling between layers

HTC Structure

(High Temperature Commensurate)




Anti-ferromagnetic coupling between layers







ITC Phase

(Intermediate Temperature Commensurate)




$$H = -J \mathbf{S}_1 \cdot \mathbf{S}_2$$

Hemert et al., PRL 51 (1983)

		J	alignment	
parallel		-		AF
linear		-		AF
T		-		AF
X		+		F
S		+		F

spin alignment: in-plane



« parallel »



$$J < 0$$




(Hemert et al.,
PRL 51 (1983)
and others)



spin alignment: inter-plane


« shifted parallel »




$$J \geq 0$$




(Hemert et al.,
PRL 51 (1983)
and others)



LTC phase




← 140 K



Jodl et al., PRB 1985

$$(\partial T / \partial P)_{\langle u^2 \rangle} = - \frac{(\partial \ln \langle u^2 \rangle / \partial P)_T}{(\partial \ln \langle u^2 \rangle / \partial T)_P}$$



$$\left. \begin{array}{l} (\partial \ln \langle u^2 \rangle / \partial T)_P = 5 \times 10^{-3} \text{ K}^{-1} \\ (\partial \ln \langle u^2 \rangle / \partial P)_T = -8 \times 10^{-2} \text{ GPa}^{-1} \end{array} \right\} (\partial T / \partial P)_{\langle u^2 \rangle} = 16 \text{ K/GPa}$$

measured: $\approx 15 \text{ K/GPa}$

Conclusions

Solid O₂ at 8 GPa is antiferromagnetically ordered at room temperature

There are three different AF structures in the δ -phase

Transitions seem to be governed by the librational motion of the O₂ molecules

Is an effect of the strong dependence of exchange interaction on the mutual orientation of neighboring molecules.

Muchas gracias por la atención!

Simposio
MALTA-Consolider
23-27 Enero 2011, Miraflores de la Sierra



www.malta-consolider.com

Moderador Mercedes Taravillo Corralo UCM

Ponencias Sistemas Moleculares I	<i>Caracterización de yunques de zafiro y moissanita a alta presión</i>	Elena del Corro García UCM
	<i>Presión interna de disoluciones acuosas</i>	Nubia Judith Mendoza Lugo UCM
	<i>Síntesis de Resinas Fósiles a alta presión. Resina, copal y ámbar</i>	Óscar Rodríguez Montoro UCM



Sistemas Moleculares I – Materiales NOCH



- Los sistemas materiales objeto de estudio específico en el desarrollo de la actividad investigadora que se propone contienen **nitrógeno, oxígeno, carbono o hidrógeno (materiales NOCH)**.
- Esta familia de materiales es extremadamente importante para nuestros propósitos por dos razones: 1) están relacionados con nuestros proyectos actuales de investigación individuales, y 2) intervienen en problemas de interés general en ciencias de la vida, de la Tierra y de los materiales.



*Pressure as a probe
on unsaturated
NOCH systems*




Sistemas Moleculares I – Materiales NOCH

Work Plan II. a.	SCIENTIFIC SCOPE	
Field Coordinator M. Taravillo	Scientific Questions	
Leading Groups UCM - UJI	- Pressure-induced polymorphism in NOCH systems - Pressure induced conformational changes in flexible molecules. Packing effects in the solid state - Non-covalent interactions and its response to pressure in the solid state - Effect of pressure on hydrogen-bonded NOCH solids. Glassy phases	
Supporting Groups All	Specific Goals	
Related Fields Water related Reactivity NOCH and minerals	- Polymorphism in sort <i>n</i>-alkanes, <i>n</i>-alcohols and benzene derivatives - Polymorphism in rigid molecules $R - C \equiv C - R'$ and $R - C \equiv N$ derivatives - Amorphous phases in alcohols and amines	

*physico-chemical phenomena
involving NOCH molecular systems*



Principales efectos de la presión en cristales moleculares



Sistemas Moleculares I – Materiales NOCH


Interés de estos estudios

Materiales sintetizados a altas p y T recuperados con propiedades tecnológicas importantes

Propiedades ópticas

Propiedades mecánicas

Superconductividad a alta temperatura





Sistemas Moleculares I – Materiales NOCH

BACKGROUND EN EL GRUPO UCM EN SISTEMAS MOLECULARES

1. Equilibrio conformacional a alta presión

1-Pentanol







2. Fenómenos de agregación a alta presión

Metanol

Agua

Agua-Metanol

Ciclohexanona







3. Transiciones de fase/polimorfismo a alta presión

Dimetilacetileno (2-butino)

Pentanol, 2-metil-2-butanol

Ciclohexano, Ciclohexanona

Teflón® (politetrafluoroetileno)





4. Reactividad Química en Fase Condensada

Monóxido de Carbono (CO)


Dimetilacetileno (2-butino)

Tetracloruro de Carbono






5. Nanomateriales bajo presión

Nanotubos de Carbono de Doble Pared



Sistemas Moleculares I – Materiales NOCH

GENERACIÓN DE CONDICIONES NO-HIDROSTÁTICAS









- Uso de yunques de **moissanita (CSi)** y de **zafiro** que permiten analizar una parte del espectro Raman de compuestos derivados del carbono que es enmascarada por el diamante.



Sistemas Moleculares I – Materiales NOCH


GENERACIÓN DE CONDICIONES NO-HIDROSTÁTICAS



Sistemas Moleculares I – Materiales NOCH


GENERACIÓN DE CONDICIONES NO-HIDROSTÁTICAS

DWCNT






GRAFITO

Desorden inducido por la presión



OTROS RESULTADOS

- High temperature microscope stage (Linkam TS1500)
 - Medidas hasta 1800 K a presión ambiente
 - Melting
 - Fonones
 - Ecuación de estado





Percy Williams Bridgman
Harvard University, USA
1882-1961
Nobel de Física 1946





Transiciones Irreversibles destacadas
-Fósforo
-CS₂

Transformaciones irreversibles de compuestos orgánicos bajo altas presiones. P. W. Bridgman y J. B. Conant, Proc. Nat. Acad. Sci. U.S. 15, 680-683 (1929).

Observaciones


- Compuestos parcialmente polimerizados al someterlos a una presión de 0.9 GPa durante 24 horas

Isopreno, 2,3-dimetil-1,3-butadieno, estireno, indeno



- Compuestos transformados al someterlos a una presión de 1.2 GPa durante 40 horas



Isobutanal, n-butanal



Sistemas Moleculares I – Materiales NOCH


Ejemplo: Benceno

Diagrama de fases



Fase a baja temperatura
y presión ambiental
Pbca

Ángulo entre los anillos de
benceno: $90^\circ 22''$



Fase a alta presión
(>1.2 GPa)
P₂₁/c

Ángulo: 120°




Sistemas Moleculares I – Materiales NOCH

REACTIVIDAD INDUCIDA POR LA PRESIÓN DEL DIMETILACETILENO





Alta presión:

Formación de Moléculas Inestables a presión
ambiente: anillos de tetrametilciclobutadieno



Descompresión:

Residuo: Material
nanográfítico




Sistemas Moleculares I – Materiales NOCH


COMPUESTOS DE INTERÉS PARA POLIMERIZAR A ALTA PRESIÓN

DITERPENOS



Abietane Type




Pimarane Type




Labdane Type



Clerodane Type




Kaurane Type




TRITERPENOS

Lupane Type




R = CH₃ Lupeol
R = CH₂OH Betulin
R = COOH Betulinic acid

Ursane Type




R = CH₃ α-Amyrin
R = COOH Ursolic acid

Oleanane Type




R = CH₃ β-Amyrin
R = COOH Oleanolic acid

Dammarane Type




Dammaradienol


TERPENOS FENÓLICOS




XII. Hinokiol



Ferruginol



XIV. 18-Norferruginol



Tatarol





Sistemas Moleculares I – Materiales NOCH

FENÓMENOS DE AGREGACIÓN A ALTA PRESIÓN

Efecto de la Presión
en el Enlace de
Hidrógeno del
Metanol Líquido






**Agregados
Metanol**




EFECTO DE LA PRESIÓN EN EL ENLACE DE HIDRÓGENO

FENÓMENOS DE AGREGACIÓN A ALTA PRESIÓN

Agua



- La presencia de agua provoca la compresión de las moléculas de metanol.
- Las variaciones con la composición se dan independientemente en agregados de agua y metanol.
- Un aumento de la presión no provoca cambios en la estructura del agua hasta que no se alcanza una composición constante de agregados de metanol.



Sistemas Moleculares I – Materiales NOCH

FENÓMENOS DE AGREGACIÓN A ALTA PRESIÓN

- Al mezclar **agua** y **metanol** el aumento de entropía que se observa es mucho más pequeño que el que se esperaría para una disolución ideal.




FIG. 3. ΔS_v for non-ionic solutes in water at 25°C. Data are from Table IV.


letters to nature

NATURE | VOL 416 | 25 APRIL 2002

Molecular segregation observed in a concentrated alcohol–water solution

S. Dixit¹, J. Crain¹, W. G. K. Poon¹, J. L. Finney² & A. K. Soper²

- Experimentos de difracción de neutrones
- Mezcla concentrada metanol-agua 7:3.
- Se concluye que las anomalías termodinámicas se deben a una mezcla incompleta a nivel molecular de las moléculas de metanol y agua.
- La estructura del agua en la mezcla es cercana a la del agua pura.



Sistemas Moleculares I – Materiales NOCH

FENÓMENOS DE AGREGACIÓN A ALTA PRESIÓN



VOLUME 91, NUMBER 15

PHYSICAL REVIEW LETTERS

week ending
10 OCTOBER 2003


Molecular Structure of Alcohol-Water Mixtures

J.-H. Guo,^{1,*} Y. Luo,^{2,†} A. Augustsson,^{1,3} S. Kashtanov,² J.-E. Rubensson,³ D. K. Shuh,⁴ H. Ågren,² and J. Nordgren³



• Espectroscopía de emisión de rayos-X.

- Para una mezcla equimolar agua-metanol, las moléculas de ambos compuestos se mezclan de forma incompleta a nivel molecular.
- Existe un nuevo mecanismo para aumentar el orden en la disolución, las moléculas de agua unen cadenas de metanol para formar anillos con 6 y 8 moléculas de metanol.






www.malta-consolider.com

Simposio

MALTA-Consolider

23-27 Enero 2011, Miraflores de la Sierra




Presión interna de disoluciones acuosas

Nubia Judith Mendoza Lugo

Departamento Química-Física


Universidad Complutense Madrid

MALTA CONSOLIDER



Contenido

1. Introducción
 2. Sistemas OH
 - Agua+Metanol
 - Agua+Etanol
 - Etanol+Metanol
 3. Características de la Experimentación
 4. Resultados
 5. Perspectivas
-




1. Introducción

Los resultados que presentaré en este trabajo forman parte de mi proyecto de tesis doctoral.

Los sistemas que estamos estudiando son: agua-metanol, agua-etanol y etanol-metanol, en función de la concentración y la presión, mediante Espectroscopía Raman.

El objetivo final de nuestro estudio es confirmar que el efecto de mezcla es similar al que ejerce la presión sobre las moléculas en estado líquido. Para ello llevaremos a cabo estudios de alta presión para los componentes puros y trataremos de correlacionar las variaciones de frecuencia, intensidad, etc. de las bandas en función de la concentración.



2. Sistemas OH

Hemos empezado nuestros experimentos con la mezcla agua-metanol, eligiendo el metanol por ser una de las más simples moléculas anfifílicas y porque poseemos resultados Raman a alta presión en ambos sistemas.

Éste estudio, junto con el del agua-etanol, sirven de referencia para acometer estudios más complejos tales como la formación de micelas, bicapas, etc en diferentes disolventes.


La mezcla etanol-metanol completa nuestro trabajo y tiene especial interés en el campo de las altas presiones.

3. Características de la Experimentación




Espectrómetro Raman BWTEK

- ✓ Excitación Láser 532nm
- ✓ Resolución $\pm 2 \text{ cm}^{-1}$
- ✓ Objetivo 10x
- ✓ Tiempo de adquisición 600-1200ms
- ✓ Acumulaciones: 30




$\text{H}_2\text{O} + \text{MeOH}$


Análisis de cuatro zonas de los espectros H₂O + MeOH




Zona de C-O



Representación de la variación con la fracción molar del MeOH





El efecto de mezcla no influye sobre las constantes de fuerza de estas vibraciones.





T. Sim. Del CO la dependencia con la concentración se puede separar en 3 zonas. Esta banda nos permitirá establecer una relación entre los efectos de solvatación y presión

Comparación de resultados

Se podría decir que cuando el H₂O hidrata una molécula anfifílica, solvata primero la mitad POLAR y después la mitad NO POLAR (CH₃).




Bending del Metilo




Observamos cómo, en este caso igual que en el bending del OCH no existe dependencia con la concentración de la frecuencia para ésta vibración.

Resonancia de Fermi




Las bandas centradas en 2845 y 2950 cm⁻¹ son las que nos van a permitir correlacionar los efectos de presión y solvatación para el gpo.CH3. Hemos comparado nuestros resultados para la banda de menor fcia con los de Dixit.






Raman Shift cm⁻¹






El efecto de mezcla se traduce en una ligera compresión del grupo CH_3 al aumentar la proporción de agua




El efecto de mezcla es un orden de magnitud mayor y de signo contrario en el enlace CO.




Zona de los OH's

Raman Shift cm⁻¹

a.s.





a.m.



Análisis de la 2da derivada de la zona de los OH's


En la posiciones de las bandas podemos ver cómo cuando pasamos de MeOH puro a fracción molar .85, la posición del mínimo se desplaza prácticamente hasta la posición de mínimo del agua pura. Indicando que pequeñas cantidades de agua alteran mucho la estructura del MeOH.








$\text{H}_2\text{O} + \text{EtOH}$

Análisis de zona de los OH's de los espectros H₂O + EtOH.




Zona de los OH's








Análisis de zona de los OH's de los espectros de EtOH+MeOH.




Zona de los OH's





5. Perspectivas

- ❑ Medir el espectro Raman de Etanol puro en función de la presión.
 - ❑ Completar el análisis de nuestros datos experimentales para las mezclas agua-etanol y etanol-metanol.
 - ❑ Hacer una interpretación de los resultados obtenidos correlacionando los efectos de presión y de hidratación de forma análoga a lo que hemos hecho con la mezcla Agua-Metanol.
 - ❑ Utilizar los resultados de la mezcla etanol-metanol para aplicaciones tanto en el campo de las altas presiones como en el estudio de sistemas biológicos.
-



Gracias por su atención...

"Síntesis de Resinas Fósiles a alta presión"

Reunión MALTA-CONSOLIDER. Enero 2011

Óscar Rodríguez Montoro

UNIVERSIDAD COMPLUTENSE DE MADRID

Departamento de Química-Física I. Universidad Complutense de Madrid

QUIM  PRES



Matter at High Pressure




Dirigido por: Valentín García Baonza y Mercedes Taravillo Corralo

OBJETIVOS


- ✓ Estudio de la influencia de la presión en la formación de resinas fósiles.
- ✓ Estudio del comportamiento de los terpenos a alta presión por métodos espectroscópicos.
- ✓ Síntesis por polimerización a alta presión de resinas fósiles. Ámbar, Copal y Resinas.
- ✓ Imitación de los procesos de fosilización por alta presión (ausencia o presencia H_3O^+).
- ✓ Establecimiento de una escala de tiempo geológica, Presión vs. Tiempo.
- ✓ ¿Búsqueda de Carbones no gráficos (combustibles fósiles)?

TERPENOS

ISOPRENO




DIVERSIDAD Y CLASIFICACIÓN



Unidades 5 C	Nombre	Total carbonos
2	Monoterpenos	10
3	Sesquiterpenos	15
4	Diterpenos	20
6	Triterpenos	30
8	Tetraterpenos	40
>20	Politerpenos	>100

NATURALEZA




CLASES Y TIPOS DE RESINAS

SECRECIONES RESINOSAS

CANALES DE RESINA




PELOS GRANULARES




TERPENOS RESINOSOS- Ej. Resina Alpina


De la oleorresina se pueden separar, su **componente sólido (colofonía o pez rubia)** y **líquido (aguarrás o esencia de trementina)**, que tienen diferentes usos en la industria.




ácido levopimarico




ácido neoabiético




ácido pimarico



ácido abiético



ácido palústrico



ácido isopimarico

TIPOS DE ÁRBOLES RESINOSOS - CONÍFERAS

Pinaceae (Pino)



Cupressaceae (ciprés y cedro)



Aracariaceae




Agathis Robusta




α -pineno




β -pineno




β -mirceno




Limoneno




Farneseno




β -felandreno




(+)-3-careno



borneol




Canfeno



Terpinoleno


FORMACIÓN DE RESINAS FÓSILES



RESINA ➡ COPAL ➡ ÁMBAR

Tiempo (millones de años)


- La **Resina** sufre un proceso de polimerización y se transforma en **Copal**, reacciones posteriores entre las cadenas de moléculas producen el **Ámbar**.
- El **Copal** es una resina del mismo origen que el Ámbar pero que va de semifósil (Pleistoceno, Mioceno) a las resinas actuales.
- El Copal tarda decenas de millones de años para cambiar a **Ámbar**. De todos los ámbares, el del Báltico es el más importante, con una presencia de ácido succínico en proporción del 3 al 8 por ciento, ha dado lugar a diferentes opiniones sobre si el auténtico ámbar debería ser sólo aquel que contuviera **ácido succínico o no**.



- Actualmente se tiende a dividir las resinas fósiles en **succinitas**, con ácido succínico, y **retinitas** sin ácido succínico, y a todas ellas si son fósiles se les considera ámbar.
- Es frecuente que algunos ámbares lleven **sulfuros (pirita) o calcita**.

- 1º pérdida de la fracción más volátil.

- Se piensa que durante la formación del ámbar, dos procesos químicos están presentes: reacciones de **condensación alquílicas** (Diels-Alder) y **polimerizaciones tipo vinílicas**.



David J. Clifford and P. G. Hatcher




Fig. 1. Polymerization of the labdatrienes communic acid (a), communal (b), and biformene (c) yielding a 14,15-polyabdatiene polymer (d).

YACIMIENTOS DE ÁMBAR (Origen geográfico)



IVº	HOLOCENO	(Actual)		
NEÓGENO	PLEISTOCENO	CALABRIENSE	VILLAFRANQUIENSE	0.01
		PLACENZIANSE	RUSCINIENSE	1.8
		ZANLAYENSE	TURQUENSE	3.4
		ESSINIENSE	VALLESENSE	5.3
		TONIENSE	ASTARAC	6.5
	MIOCENO	RAVALLIENSE	PROCONENSE	11
		LANGHIENSE	RAMBLIENSE	14.5
		BURDIGALIENSE	AGENIENSE	16
		AQUITANIENSE	ARVERNIENSE	23.5
		SESSIENSE	SUEVIENSE	28
PALEÓGENO	OLIGOCENO	SESSIENSE	PERMANENSE	34
		BARONIENSE	SUP. MEDIO INF.	37
	EOCENO	LUTICIENSE	NEUSTRIENSE	40
		YPRESIENSE	CUISIENSE	46
		THANIENSE	ILERIENSE	53
CRETÁCICO	PALEOCENO	DANIENSE	GARUMN	59
		MAASTRICHTIENSE		65
	SUPERIOR	CRANIANSE		72
		SENNANENSE		83
		TURONIENSE		87
	INFERIOR	CENOMANIENSE	UTRILLAS	91
		ALBIENSE		96
		APTIENSE	URGONTIANA	108
		BARREMIENSE		114
		HAUTERVIENSE	WEALD	116
		VALANGIENSE		122
		BERRIASIENSE	PURBECK	130
				135

- El **Ámbar de Canadá** (Chemawinita o Cedarita) es de color marrón. Es muy antiguo, se cree procede de coníferas del Cretáceo (60 a 130 millones de años). Es también retinita.
- El **Ámbar de España** puede ser de yacimientos del Cretácico en Álava, Teruel y Cantabria, este último (cerca de **la cueva de El Soplao**) descubierto en 2007, de alrededor de hace 100 millones de años y de gran interés por la cantidad de insectos y artrópodos (arañas) incluidos. Los **otros yacimientos** de Ámbar del Cretácico tanto en España como en el resto de los países donde se han encontrado (Líbano, Israel, Jordania, oeste de Francia, sur de Gran Bretaña, Noreste de España) no tienen tantos insectos. Se ha encontrado también un raro Ámbar azul de este período, las mejores piezas son las que se formaron de forma subaérea (en cavidades dentro del tronco).
- El **Ámbar del Báltico o Succino**, de los primeros yacimientos conocidos y de los más importantes. Procede de las costas de Dinamarca, Suecia, norte de Alemania, Polonia y parte de Rusia. La mayor parte se recoge del mar, flotando o en el fondo y en menor proporción en excavaciones o minas. Suele ser de color amarillo miel a pardo, a veces con tonos rojizos anaranjados y hasta casi blancos, transparentes o nebulosos. Procede de coníferas del Eoceno (40 a 60 millones de años). Es el de mayor proporción de ácido succínico, del 3 al 8 por ciento, por tanto es succinita. Su uso y comercio se remonta hasta la prehistoria.
- El **Ámbar de Myanmar (antigua Birmania)**, llamado Burmita, es de colores fuertes, principalmente marrón oscuro y algo turbios. Es succinita por llevar un 2 por ciento de ácido succínico. Muy fluorescente. Lleva frecuentemente calcita en el interior de las fisuras y le da una apariencia moteada característica. Procede de burseráceas del Eoceno (40 a 60 millones de años).
- El **Copal de Rumanía (Rumanita)** es muy rico en colores: pardo amarillento a marrón profundo, granate, verdoso y casi negro. Lleva gran cantidad de sulfuros que le proporcionan el color característico. La proporción de ácido succínico es del 1 al 5 por ciento, es succinita. Se originó en el Mioceno a partir de leguminosas (10 a 25 millones de años).
- El **Copal de Sicilia (Simetita)**, nombre derivado de su lugar de origen, el río Simeto, es uno de los más bellos. De color rojo oscuro, azulado, verdoso, y raramente amarillo. Suele ser transparente. No lleva ácido succínico, es retinita; pero tiene gran cantidad de sulfuros. Procede de burseráceas del Mioceno (10 a 25 millones de años).
- El **Copal de México**, se presenta en el Distrito de Chiapas. Tonos de rojo, amarillo y casi negro. Es una retinita procedente de leguminosas (Hymenaea) del Mioceno.
- El **Copal de la República Dominicana** es de color amarillo pálido a marrón, raramente rojo, a veces azul, que es el más apreciado. Muy fluorescente a la luz UV. Procede de leguminosas tipo algarrobo, del Mioceno (10 a 25 millones de años). No tiene ácido succínico, es retinita.
- **Otros yacimientos más jóvenes** se encuentran también en EE.UU., Japón, Tanzania (Copal joven procedente del Pleistoceno, 1 a 10 millones de años), y en Nueva Zelanda (Ambrita).

ESTUDIOS PREVIOS DE TERPENOS A ALTA PRESIÓN

(...) Isoprene is polymerized to rubber (...)




Conant, J. B., Tongberg, C. O. Polymerization reactions under high pressure. I. Some experiments with isoprene and butyraldehyde. *J. Am. Chem. Soc.* 52, 1659 (1930).

> 1,2 GPa

Studies in the Terpene Series. XVII. The Thermal Isomerization of Pinane at High Pressure

Ipatieff et al. Studies in the Terpene Series. XVII. The Thermal Isomerization of Pinane at High Pressure. *J. Am. Chem. Soc.* (1953)




α -pineno β -pineno


400 °C; 80 MPa

Organic Reactions Under High Pressure. IV. The Dimerization of Isoprene

C. Walling and J. Peisach. Organic Reactions Under High Pressure. IV. The Dimerization of Isoprene. *J. Am. Chem. Soc.* (1958)




> 1,2 GPa



Dimerization and Polymerization of Isoprene at High Pressures

Citroni, M; Ceppatelli, M; Bini, R, et al. Dimerization and Polymerization of Isoprene at High Pressures. *J. Phys. Chem. B* 2007, 111, 3910-3917



< 2,6 GPa

Figure 1. IR spectra recorded during the room temperature reaction of isoprene at 1.15 GPa. The first spectrum (0 h) of the kinetics is entirely due to the monomer and is measured just after the pressure of the experiment has been set. The arrows indicate the absorption bands of the product that do not overlap with monomer peaks.





Figure 12. Possible dimerization paths leading, from top to bottom, to sylvestrene (a), limonene (b), 4-ethenyl-2,4-dimethylcyclohexene (c), and 4-ethenyl-1,4-dimethylcyclohexene (d).

ESPECTROSCOPIA DEL ÁMBAR (I)

TRANSFORMACIONES MOLECULARES COMUNES DEPENDIENTES DEL TIEMPO ACAECIDAS EN LAS RESINAS FÓSILES


- (1) Pérdida progresiva de dobles enlaces olefínicos (especialmente los enlaces exocíclicos).
- (2) Desaparición de grupos funcionales (oxidación extrema).
- (3) Incremento proporcional de componentes aromatzados, pero muy lejos en proporción con los porcentajes encontrados en maderas fosilizadas o incluso en bancos de carbón.

"Spectroscopic Investigations of Amber". C. W. Beck. Applied Spectroscopy Reviews, 1520-569X, Volume 22, Issue 1, 1986, Pages 57-110



ESPECTROSCOPIA DEL ÁMBAR (II)

“Fourier transform-Raman spectroscopy of amber”. H. G. M. Edwards, D. W. Farwell. Spectrochimica Acta Part A 52 (1996) 1119-1125




Asignaciones vibracionales del ámbar

Table 1
Wavenumbers/cm⁻¹ and vibrational assignments for amber

ν/cm^{-1}	Approximate assignment of vibrational mode
2965 m, sh	$\nu(\text{CH}_3)$ stretching
2930 s	$\nu(\text{CH}_2)$ stretching
2892 m, sh	$\nu(\text{CH}_2)$ stretching
2871 ms	$\nu(\text{CH}_2)$ stretching
2850 m, sh	$\nu(\text{CH}_2)$ stretching
2728 w	$\nu(\text{CH}_2-\text{CH}_2)$ terminal
1734 mw	$\nu(\text{C=O})$ stretching ester
1646 ms	$\nu(\text{C=C})$ stretching
1610 m, w, sh	$\nu(\text{C=C})$ stretching
1450 s	$\delta(\text{CH}_3)$ deformation
1420 w	$\delta(\text{CH}_3)$ deformation
1356 m	$\delta(\text{CH}_2)$ twisting
1298 m	$\delta(\text{CH}_2)$ twisting
1245 w	
1205 m	
1143 w	
1108 w	$\nu(\text{C}-\text{C})$
1060 w	$\nu(\text{C}-\text{Cl})$; $\nu(\text{C}-\text{O})$
1034 w	$\nu(\text{C}-\text{O})$
1014 vw	$\nu(\text{C}-\text{O})$
976 m, br	$\rho(\text{CH}_2)$
938 mw, br	$\rho(\text{CH}_2)$
881 mw	$\nu(\text{COC})$
838 mw, br	
742 w	
718 m	
696 w	
651 mw	$\nu(\text{C}-\text{S})$ stretching
540 w	$\nu(\text{COC})$ in-plane deformation
504 m, br	$\nu(\text{S}-\text{S})$ stretching; $\nu(\text{COC})$
454 m, br	$\delta(\text{CCO})$
375 w	$\delta(\text{CCC})$
310 w	$\tau(\text{CCO})$

Se comparan ámbares de diferentes tipos, que aunque tienen un espectro Raman similar en la región de 2000-3000 cm⁻¹, al compararlos con los de resina de pino contemporánea, se observa a 3020 cm⁻¹ las bandas de alargamiento $\nu(\text{CH}$, stretching), debidas al enlace del grupo C=CH- que está ausente en las muestras de resinas fosilizadas.




ÁMBAR DE LA CUEVA DEL SOPLAO-Patrimonio

“Terpenoids in extracts of Lower Cretaceous ambers from the Basque-Cantabrian Basin (El Soplao, Cantabria, Spain): Paleochemotaxonomic aspects”. C. Menor-Salván, M.Najarro, F. Velasco, I.Rosales, F. Tornos, Bernd R. T. Simoneit. Organic Geochemistry xxx (2010) xxx-xxx. In Press.


2 tipos de ámbar de diferente origen biológico

Terpenoides fenólicos:

- Ferruginol, Totarol y Hinokiol
- Ácidos pimárico e isopimárico
- Azulenos



- Ácidos pimárico e isopimárico
- Ausencia de terpenos fenólicos




El FTIR de ambos tipos de ámbar encontrados tiene espectros similares.

COMPUESTOS DE INTERÉS PARA POLIMERIZAR A A. P.


DITERPENOS

Abietane Type




Abietic acid

Pimarane Type




Pimaric acid

Labdane Type




Communic acid




Ozoic acid

Clerodane Type



Hardwickiic acid


Kaurane Type



Trachylobanic acid


TRITERPENOS

Lupane Type




R = CH₃ Lupeol
R = CH₂OH Betulin
R = COOH Betulinic acid

Ursane Type




R = CH₃ α-Amyrin
R = COOH Ursolic acid

Oleanane Type




R = CH₃ β-Amyrin
R = COOH Oleanolic acid

Dammarane Type




Dammaradienol


TERPENOS FENÓLICOS




XII. Hinokiol



Ferruginol



XIV. 18-Norferruginol





Totarol


ESTRATEGIAS PARA LOS EXPERIMENTOS

REACCIONES DIELDS-ALDER Y VINÍLICAS

Polymeric Skeletons



IV ^o	HOLOCENO	(Actual)	0.01
	PLEISTOCENO	CALABRENSIS	1.8
NEÓGENO	PLIOCENO	PLACIENSIS	3.4
		ZANCLAYENSIS	5.3
	MIOCENO	MESSENIENSIS	6.5
		TORTONENSIS	11
		SERRAVALLENENSIS	14.5
PALEÓGENO	OLIGOCENO	LANGHEIS	16
		BURDIGALENSIS	20
		AQUITANENSIS	23.5
		CHATTIENSIS	28
		RUPELIENSIS	34
	EOCENO	PRABOHENSIS	37
		BARTONENSIS	40
	PALEOCENO	LUTICIENSIS	46
		YPRASIENSIS	53
		THAMETIENSIS	59
CRETÁCICO	SUPERIOR	DANIESENSIS	65
		MAASTRICHTIENSIS	72
		CAMPANENSIS	83
		SANTONENSIS	87
		CONIAGENSIS	91
	INFERIOR	TURONIENSIS	96
		CENOMANENSIS	108
		ALBIENSIS	114
		APTIENSIS	116
		BARREMENSIS	122



- ¿Qué λ del láser usar?
- ¿Hasta qué presiones llegar?
- ¿Qué tipo de yunques usar?
- ¿Evitar la luminiscencia o realizar medidas de ellas?
- ¿Cómo utilizar el ámbar sin ir en contra del Código Penal?



“gracias por la atención”

QUIM  PRES



 CONSOLIDER
ALTA

Matter at High Pressure

PROGRAMA
ingenio
2010

Óscar Rodríguez Montoro

UNIVERSIDAD COMPLUTENSE DE MADRID

Departamento de Química-Física I. Universidad Complutense de Madrid

Funciones de Distribución del Número de Electrones (EDF): Conexión con los orbitales del hueco de Fermi promediado en un dominio del espacio real (DAFH)

The Journal of Chemical Physics **131**,124125 (2009)

E. Francisco, A. Martín Pendás y M. A. Blanco

Universidad de Oviedo

Contenido

- 1 Introducción
- 2 Metodología
- 3 Resultados y discusión
- 4 Resumen y conclusiones

I. Introducción

- La obtención de información química relevante a partir de Ψ
PROBLEMA NO TRIVIAL. La dificultad $\uparrow$ al $\uparrow$ la complejidad de Ψ .
- De entre los muchos métodos desarrollados ...

- ▶ **Funciones de Distribución del Número de Electrones (EDF)**

Si $\Omega_1 + \Omega_2 + \dots + \Omega_m = \mathcal{R}^3$, $\Omega_k \cap \Omega_{l \neq k} = \emptyset$, $n_1 + n_2 + \dots + n_m = N$

EDF = $\{P(n_1, n_2, \dots, n_m)\} = \{P(S)\}$ donde

$P(S)$ = Probabilidad de hallar n_1 e⁻ en Ω_1 , n_2 en $\Omega_2, \dots$, n_m en Ω_m .

Las EDFs son útiles para ...

- ★ Aditividad aproximada de las propiedades de grupos funcionales.
 - ★ Particiones energéticas.
 - ★ Definir nuevos conceptos (e.g. Est. resonantes en el espacio real, RSRS).
 - ★ Clasificación de enlaces químicos
 - ★ Caminos de enlace $\equiv$ canales privilegiados de intercambio mecano-cuántico.
- ▶ **Domain Averaged Fermi Hole (DAFH) analysis (Robert Ponec)**
 $\rho^{\text{hole}}(\mathbf{r}|\mathbf{r}_1) \rightarrow \langle \mathbf{r}_1 \rangle_{\Omega} \rightarrow G^{\Omega}(\mathbf{r}) \Rightarrow$
Muy útil para entender/interpretar diferentes aspectos de la naturaleza del enlace químico, e.g.
 - ★ Moléculas hipervalentes y/o ricas en electrones.
 - ★ Moléculas recién sintetizadas.
 - ★ Estructura de MOX_4 (M=Os, Ru; X=F, Cl, Br, I).
 - ★ Enlace metal-metal en $\text{Re}_2\text{Cl}_8^{2-} \dots$

I. Introducción

Objetivos de este trabajo:

- Establecer la conexión entre EDF & DAFH.
- Análisis del enlace en ejemplos sencillos.

Condiciones:

- $\Psi(1, N) = \det [\varphi_1(1)\varphi_2(2) \dots \varphi_N(N)]$.
- $\mathcal{R}^3 = \Omega + \Omega'$ (QTAIM)

II. Metodología: A. DAFH

- $\rho^{\text{hole}}(\mathbf{r}_2|\mathbf{r}_1) = \rho(\mathbf{r}_2) - \frac{\rho_2(\mathbf{r}_1, \mathbf{r}_2)}{\rho(\mathbf{r}_1)} \Rightarrow \rho_{\Omega}^{\text{hole}}(\mathbf{r}_2) = \rho(\mathbf{r}_2) - \frac{\int_{\Omega} \rho_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1}{\int_{\Omega} \rho(\mathbf{r}_1) d\mathbf{r}_1}$

$$\text{DAFH} \equiv G^{\Omega}(\mathbf{r}) = N_{\Omega} \rho_{\Omega}^{\text{hole}}(\mathbf{r}) = \int_{\Omega} \rho_2^{\text{xc}}(\mathbf{r}_1, \mathbf{r}) d\mathbf{r}_1 = \sum_{i,j}^N \varphi_i(\mathbf{r}) \langle \varphi_i | \varphi_j \rangle_{\Omega} \varphi_j^*(\mathbf{r}).$$

- $G^{\Omega}(\mathbf{r}) = \sum_{i,j}^N \varphi_i(\mathbf{r}) \mathbf{S}_{ij} \varphi_j^*(\mathbf{r}) = \boldsymbol{\varphi} \mathbf{S} \boldsymbol{\varphi}^{\dagger}$, donde $\mathbf{S} \equiv \langle \boldsymbol{\varphi} | \boldsymbol{\varphi} \rangle_{\Omega}$.

- $G^{\Omega}(\mathbf{r})$ tiene muchas propiedades importantes.

- $\text{Diag } \mathbf{S} \Rightarrow \mathbf{U} \text{ y } \mathbf{n} \Rightarrow \text{DAFH MOs } \boldsymbol{\phi} = \boldsymbol{\varphi} \mathbf{U} \text{ \& } G^{\Omega}(\mathbf{r}) = \sum_{i=1}^N n_i |\phi_i(\mathbf{r})|^2.$

- $\langle \phi_i | \phi_j \rangle_{\mathcal{R}^3} = \delta_{ij} \quad \langle \phi_i | \phi_j \rangle_{\Omega} = \delta_{ij} n_i \quad (\star) \quad \langle \phi_i | \phi_j \rangle_{\Omega'} = \delta_{ij} (1 - n_i).$

- $\Psi(1, N)$ es invariante frente a $\mathbf{U} \Rightarrow$

$$\Psi(1, N) = \det [\varphi_1(1) \varphi_2(2) \dots \varphi_N(N)] = \det [\phi_1(1) \phi_2(2) \dots \phi_N(N)].$$

- $\rho(\mathbf{r}) = \sum_i^N |\phi_i(\mathbf{r})|^2$, $\parallel$ Cada ϕ_i contribuye con un e^- a ρ !!

- A partir de $(\star) \Rightarrow n_i = \text{Probabilidad de hallar el } e^- \text{ descrito con } \phi_i \text{ en } \Omega$
 $= \text{Fracción del } e^- \text{ descrito con } \phi_i \text{ contenido en } \Omega$

II. Metodología: B. EDF

- La probabilidad de que exactamente n electrones estén en Ω y $N - n$ en Ω' es:

$$\begin{aligned} p^\Omega(n) &= \frac{N!}{n!(N-n)!} \int_{\Omega} d\mathbf{r}_1 \cdots d\mathbf{r}_n \int_{\Omega'} d\mathbf{r}_{n+1} \cdots d\mathbf{r}_N |\Psi|^2 \\ &= \frac{1}{n!(N-n)!} \sum_{q \in \text{perm}(1, \dots, N)} \det[\tilde{\mathbf{S}}_{ij}(q_j)], \end{aligned} \quad (1)$$

donde $\tilde{\mathbf{S}}_{ij}(q_j) = \begin{cases} \langle \phi_i | \phi_j \rangle_{\Omega} = \delta_{ij} n_i & 1 \leq q_j \leq n \\ \langle \phi_i | \phi_j \rangle_{\Omega'} = \delta_{ij} (1 - n_i) & n+1 \leq q_j \leq N \end{cases}$

- ¡¡ $\tilde{\mathbf{S}}$ es diagonal en la base DAFH !!

- Si $\mathbf{p}_1^i = (n_i, 1 - n_i) \Rightarrow \{p^\Omega(n), n = 0, 1, \dots, N\} \equiv \mathbf{p}_N = \otimes_{i=1}^N \mathbf{p}_1^i$.

CONCLUSIÓN: Si $\Psi = \text{Det}(\text{Slater}) \Rightarrow$ Los electrones son estadísticamente independientes cuando son descritos con los MO's que diagonalizan el Huevo de Fermi.

II. Metodología

EJEMPLO ($N = 3 \Rightarrow 4$ posibilidades)

$$\otimes_{i=1}^3 p_1^i = (n_1, 1 - n_1) \otimes (n_2, 1 - n_2) \otimes (n_3, 1 - n_3) = (p^\Omega(3), p^\Omega(2), p^\Omega(1), p^\Omega(0)):$$

$$p^\Omega(3) = n_1 n_2 n_3$$

$$p^\Omega(2) = n_1 n_2 (1 - n_3) + n_1 (1 - n_2) n_3 + (1 - n_1) n_2 n_3$$

$$p^\Omega(1) = n_1 (1 - n_2) (1 - n_3) + (1 - n_1) n_2 (1 - n_3) + (1 - n_1) (1 - n_2) n_3$$

$$p^\Omega(0) = (1 - n_1) (1 - n_2) (1 - n_3)$$

- $p^\Omega(3)$ = Producto de 3 eventos independientes.
- $p^\Omega(2)$ = Suma de 3 productos de 3 eventos independientes.
- $p^\Omega(1)$ = Suma de 3 productos de 3 eventos independientes.
- $p^\Omega(0)$ = Probabilidad de 3 eventos independientes.

Obviamente: Cada $p^\Omega(n)$ es invariante frente al intercambio de 2 electrones cualesquiera.

II. Metodología: Indicadores químicos del enlace

- Índice de deslocalización (DI) Ω y $\Omega' \sim$ Orden de enlace Ω y Ω'

$$\delta^{\Omega, \Omega'} = \sum_{i=1}^N 2n_i(1 - n_i) = \sum_{i=1}^N \delta_i^{\Omega, \Omega'}$$

- ▶ Los electrones contribuyen de manera independiente al enlace.
- ▶ ϕ_i muy localizado en Ω ($n_i \simeq 1$) u Ω' ($n_i \simeq 0$) $\Rightarrow \delta_i \simeq 0$ (CORE).
- ▶ Deslocalización apreciable en Ω y Ω' ($0 < n_i < 1$) $\Rightarrow \delta_i \neq 0$ (VALENCIA).
- ▶ Igual deslocalización de ϕ_i en Ω y Ω' ($n_i \simeq 0.5$) $\Rightarrow \delta_i^{\max} \simeq 0.5$.

- En capas cerradas cada ϕ_i está duplicado $\Rightarrow$

$$\delta^{\Omega, \Omega'} = \sum_{i=1}^{N/2} 4n_i(1 - n_i) = \sum_{i=1}^{N/2} \delta_i^{\Omega, \Omega'}$$

El máximo valor posible de cada $\delta_i^{\Omega, \Omega'}$ es entonces 1.0.

III. Resultados y discusión

CASOS SENCILLOS

- Bien descritos con Ψ 's monodeterminantes.
- Diagramas de orbitales moleculares LCAO conocidos.

SISTEMAS

- Moléculas diatómicas: $\text{H}_2, \text{He}_2, \text{Li}_2, \text{Be}_2, (\text{B}_2, {}^3\Sigma_u^+), \text{C}_2, \text{N}_2, (\text{O}_2, {}^3\Sigma_g^-)$ y F_2 .
- Hidruros: AH_n ($\text{A}=\text{Li}, \text{Be}, \text{B}, \text{C}, \text{N}, \text{O}, \text{F}$)


III. Resultados y discusión

molécula H_2

- $S^\alpha = S^\beta = \langle \sigma_g | \sigma_g \rangle_\Omega = 0.5$
- $(\phi_1^\alpha = \phi_1^\beta = \sigma_g)$ y $(n_1^\alpha = n_1^\beta = 0.5)$
- $(\delta_i^\alpha = \delta_i^\beta = 0.5)$



- El e^- α no siente al β y *viceversa*
- En general, $\Psi = \text{Det de Slater} \Rightarrow$
Bloques α y β son indep.



molécula He_2

- En este caso $\Rightarrow S^\alpha = S^\beta =$

$$\begin{pmatrix} \langle \sigma_g | \sigma_g \rangle_\Omega & \langle \sigma_g | \sigma_u \rangle_\Omega \\ \langle \sigma_u | \sigma_g \rangle_\Omega & \langle \sigma_u | \sigma_u \rangle_\Omega \end{pmatrix}$$

- con $\langle \sigma_g | \sigma_g \rangle_\Omega = \langle \sigma_u | \sigma_u \rangle_\Omega = 0.5$ y
 $\langle \sigma_g | \sigma_u \rangle_\Omega = \langle \sigma_u | \sigma_g \rangle_\Omega \approx 0.5$.
- $n_1 \approx 0.0, n_2 \approx 1.0$
 $\Rightarrow \phi_1 = 1\sigma_g + 1\sigma_u = 1s_A$
 $\Rightarrow \phi_2 = 1\sigma_g - 1\sigma_u = 1s_B,$
 $\delta_1 = \delta_2 \approx 0.0$


III. Resultados y discusión

molécula Li_2

- $(3 \times 3)\mathbf{S}^\alpha = \mathbf{S}^\beta, 1\sigma_g, 1\sigma_u, 2\sigma_g$.

$$\mathbf{S}^\alpha = \left(\begin{array}{cc|c} 0.5 & 0.5 & 0.0 \\ 0.5 & 0.5 & \simeq 0.0 \\ \hline 0.0 & \simeq 0.0 & 0.5 \end{array} \right)$$

- | | 1 | 2 | 3 |
|------------|----------------|----------------|---------------------|
| n_i | 1.000 | 0.000 | 0.500 |
| ϕ_i | $\approx 1s_A$ | $\approx 1s_B$ | $\approx 2\sigma_g$ |
| δ_i | 0.000 | 0.000 | 0.500 |
- 2 MO's son simétricos y casi completamente localizados en ambos núcleos de Li, y 1 MO deslocalizado por igual en ambos núcleos, $\delta \approx 1.0$.



III. Resultados y discusión

- **molécula Be₂: Cualitativamente muy similar a He₂**
 - ▶ $(S^\alpha = S^\beta = 4 \times 4): (1\sigma_g, 1\sigma_u) \rightarrow (1s_\Omega, 1s_{\Omega'})$ y $(2\sigma_g, 2\sigma_u) \rightarrow (2s_\Omega, 2s_{\Omega'})$.
- DAFH HF: **H₂, Li₂ = Enlace sencillo, He₂, Be₂ = Enlace residual.**
- **molécula B₂: Capa abierta: $(S^\alpha = 6 \times 6, S^\beta = 4 \times 4)$.**

-

	$n_1(\delta_1)$	$n_2(\delta_2)$	$n_3(\delta_3)$	$n_4(\delta_4)$	$n_5(\delta_5)$	$n_6(\delta_6)$
α	0.0 (0.0)	1.0 (0.0)	0.16 (0.26)	0.84 (0.26)	0.5 (0.5)	0.5 (0.5)
β	0.0 (0.0)	1.0 (0.0)	0.16 (0.26)	0.84 (0.26)		

- ▶ $\phi_{1,2} \simeq 1\sigma_g \pm 1\sigma_u \simeq (1s_\Omega, 1s_{\Omega'})$ $\phi_{3,4} \simeq 2\sigma_g \pm 2\sigma_u + \epsilon(1\sigma_g \pm 1\sigma_u)$ (α y β).
- ▶ $(\phi_5, \phi_6) = (1\pi_{ux}, 1\pi_{uv})$ Orbitales canónicos de HF

- molécula C_2 ($S^\alpha = S^\beta = 6 \times 6$)

- Similar a B_2 añadiendo 2 e^- $\beta \Rightarrow (\phi_5, \phi_6) = (1\pi_{ux}, 1\pi_{uy}) \Rightarrow \delta \uparrow$ respecto a B_2 .



- molécula N_2 ($S^\alpha = S^\beta = 7 \times 7$)

$n_1(\delta_1)$	$n_2(\delta_2)$	$n_3(\delta_3)$	$n_4(\delta_4)$	$n_5(\delta_5)$	$n_6 = n_7$
0.0 (0.0)	1.0 (0.0)	0.01 (0.01)	0.99 (0.01)	0.5 (0.5)	0.5

- $\phi_5 \simeq c_1 2\sigma_g + c_2 3\sigma_g$ (ϕ_6, ϕ_7) = ($1\pi_{ux}, 1\pi_{uy}$) orbitales canónicos de HF,
 $\delta = 3.0$

III. Resultados y discusión

molécula O_2 (capa abierta) ($S^\alpha = 9 \times 9$, $S^\beta = 7 \times 7$).



- $\beta \simeq N_2 \Rightarrow 3 e^-$
(uno σ y 2 π) con
máxima deslocalización.
($n_i = 0.5$).
- $\alpha: \pi_{ux}^1 \pi_{gx}^1 \pi_{uy}^1 \pi_{gy}^1$
Rep. Pauli entre
(π_{ux}, π_{gx}) y (π_{uy}, π_{gy})
 $\Rightarrow$ Localización.


III. Resultados y discusión

¿Cómo recuperar la EDF a partir de los n_i ?

- Si $n_i \simeq 1.0 \Rightarrow \phi_i \in \text{CORE de } \Omega$ o si $n_i \simeq 0.0 \Rightarrow \phi_i \in \text{CORE de } \Omega'$, y pueden excluirse del cálculo de la EDF

$$\Rightarrow \mathbf{p}_N = \bigotimes_{i \in \text{deslocalizado}} \mathbf{p}_1^i.$$

- EDF completa = Producto directo de las $\mathbf{p}_1^i$'s de valencia.
- Ejemplo, molécula O_2



- aprox EDF $\alpha = \mathbf{p}_1^5 = (0.5, 0.5)$
- aprox EDF $\beta = \mathbf{p}_1^5 \otimes \mathbf{p}_1^6 \otimes \mathbf{p}_1^7 = (0.5, 0.5) \otimes (0.5, 0.5) \otimes (0.5, 0.5) = (0, 0, 0.125, 0.375, 0.375, 0.125, 0, 0)$

III. Resultados y discusión

Hidruros AH_n ($A=Li, Be, B, C, N, O, F$) (Capa cerrada)

- Diferentes definiciones de Ω y Ω' (excepto en LiH y FH),
- Nuestra elección $\Omega = H$ y $\Omega' = AH_{n-1}$.
- Resultado: Todos los ϕ_i excepto ϕ_1 tienen $n_i \approx 0.0$
 $\Rightarrow \phi_2, \phi_3, \dots$ localizados en AH_{n-1} .
- Sólo el par (α, β) asociado a ϕ_1 está deslocalizado en Ω y Ω' .

	LiH	BeH ₂	BH ₃	CH ₄	NH ₃	H ₂ O	HF
n_1	0.9490	0.9129	0.8043	0.4890	0.3057	0.1880	0.1186
δ_1	0.0967	0.1591	0.3148	0.4998	0.4245	0.3053	0.2091
δ	0.1040	0.1859	0.4003	0.5560	0.4622	0.3278	0.2227

- El grado de localización de ϕ_1 en H decrece del LiH al HF:
 $\Rightarrow$ IMAGEN DAFH de enlace covalente polarizado sencillo entre Ω y Ω' .

IV. Resumen y conclusiones

- DAFH MO's para $\Psi(1, N)$ Det Slater $\Rightarrow$ Electrones estadísticamente independientes: $\{p^\Omega(n)\} \equiv \mathbf{p}_N = \otimes_{i=1}^N \mathbf{p}_1^i$.
- $n_i \neq$ Ocupación de ϕ_i sino que es la probabilidad de que un electrón descrito con ϕ_i esté en la región Ω
- DAFH MO's con $n_i \simeq 1$ ó $n_i \simeq 0$ están muy localizados en Ω u Ω' y no contribuyen al enlace, $\delta_i \simeq 0.0$, mientras que los MO's máximamente enlazantes ($\delta_i \simeq 0.5$) son los más deslocalizados en Ω y Ω' simultáneamente, $n_i \simeq 0.5$.


¿ Validez y posible extensión a $\Psi(1, N)$ Multi-Det ?


Magnetism in solid oxygen under pressure

Stefan Klotz

IMPMC

Université P&M Curie, Paris








A.L. Lavoisier
(1743-1794)


ὀξύς (oxys) -γενής (-genēs)

« acid former »

Oxygen






AO



AO


1848: Faraday:	Paramagnetic nature of O ₂
1910: Kamerlingh Onnes:	Magn. suscept. of liquid and solid O ₂
1963: Alikhanov:	Magn. structure of solid O ₂ (neutrons, 1 bar)
2004: Goncharenko:	Mag. struct. of δ -O ₂ under pressure (neutrons, 6 GPa)







O₂ at 1 bar

β : short-range ordered





α : antiferromagnetic



α - oxygen







$$b/a = 0.79 \neq 3^{1/2}/2$$




δ - oxygen



Nuclear Structures





Previous high P neutron experiments




Goncharenko, PRL 2004



Infrared absorption (diamond anvil cell)



Gorelli et al., PRB 2008



Sample:
 80 mm^3

TiZr gasket


cBN

$\sim 10 \text{ GPa}$ for $V = 100 \text{ mm}^3$

$\sim 30 \text{ GPa}$ for $V = 10 \text{ mm}^3$

3.5-320 K

see also talk Domingo Martinez
martes 17:40





D20 high intensity diffractometer
ILL, Grenoble, France

Future:

XTREMED: dedicated high P diffractometer at ILL
(Javier Campo Ruiz miércoles 10:45)

Sample loading



Condensation at 77 K

Gas loader






$P_{\max} = 2000 \text{ bar}$
Compatible with hydrogen



Difficulties

- Texture (imperfect powder) due to the formation of single crystals
- Strong compressibility: $\sim 50\%$ from 1-8 GPa
- Uncontrollable download
- Analysis: anisotropic magnetic form factor







220 K



145 K

100 K



LTC Structure



(Low Temperature Commensurate)




Ferromagnetic coupling between layers

HTC Structure

(High Temperature Commensurate)




Anti-ferromagnetic coupling between layers







ITC Phase

(Intermediate Temperature Commensurate)




$$H = -J \mathbf{S}_1 \cdot \mathbf{S}_2$$

Hemert et al., PRL 51 (1983)

		J	alignment	
parallel		-		AF
linear		-		AF
T		-		AF
X		+		F
S		+		F

spin alignment: in-plane



« parallel »



$$J < 0$$




(Hemert et al.,
PRL 51 (1983)
and others)



spin alignment: inter-plane


« shifted parallel »




$$J \geq 0$$




(Hemert et al.,
PRL 51 (1983)
and others)



LTC phase




← 140 K



Jodl et al., PRB 1985

$$(\partial T / \partial P)_{\langle u^2 \rangle} = - \frac{(\partial \ln \langle u^2 \rangle / \partial P)_T}{(\partial \ln \langle u^2 \rangle / \partial T)_P}$$



$$\left. \begin{array}{l} (\partial \ln \langle u^2 \rangle / \partial T)_P = 5 \times 10^{-3} \text{ K}^{-1} \\ (\partial \ln \langle u^2 \rangle / \partial P)_T = -8 \times 10^{-2} \text{ GPa}^{-1} \end{array} \right\} (\partial T / \partial P)_{\langle u^2 \rangle} = 16 \text{ K/GPa}$$

measured: $\approx 15 \text{ K/GPa}$

Conclusions

Solid O₂ at 8 GPa is antiferromagnetically ordered at room temperature

There are three different AF structures in the δ -phase


Transitions seem to be governed by the librational motion of the O₂ molecules

Is an effect of the strong dependence of exchange interaction on the mutual orientation of neighboring molecules.

Muchas gracias por la atención!

Topological partition of the elastic constants of crystals into atomic contributions

Alberto Otero-de-la-Roza and Víctor Luaña



Departamento de Química Física y Analítica, Universidad de Oviedo


48 EHPRG International Conference, Uppsala, 25–29 July 2010



- 1 Quantum Theory of Atoms in Molecules (QTAIM) partition of the space
- 2 Partition of the bulk modulus B and $\kappa = B^{-1}$
 - Influences on B_{Ω} and κ_{Ω}
- 3 Elastic equation of state
- 4 How to partition the c_{ij} and s_{ij} elastic constants
 - Partitioning c_{11}
 - Partitioning c_{12}
 - Partitioning s_{11} and s_{12}
 - Partitioning c_{44} and s_{44}
 - Internal consistency of the partitioning
- 5 Partitioning in four families of oxide crystals
- 6 Conclusions
- 7 Acknowledgement



Quantum Theory of Atoms in Molecules (QTAIM) partition of the space




C₂H₄: ethylene

- 3D space is divided into *atomic basins* by the $\nabla\rho(\mathbf{r})$ gradient vector field.
- Each basin is such that $\nabla\rho(\mathbf{r}_s) \cdot \mathbf{n}(\mathbf{r}_s) = 0$ (zero flux condition), i.e. gradient lines do not cross the *inter-atomic surface* (IAS) that separates two bonded basins.
- The only exception is a single *bond path*, the crest or ridge that connects two bonded nuclear peaks and crosses the IAS through the *bond critical point* or mountain pass.

- The zero flux condition guarantees that all quantum mechanical operators are locally well defined, hermitian, and can be integrated on each basin to produce the atomic contribution to the properties of the whole system: $\langle F \rangle_{\Omega} = \int_{\Omega} \hat{F}[\rho] \rho(\mathbf{r}) d\mathbf{r}$.
- Basin contributions are strictly additive, $\langle F \rangle = \sum_{\Omega} \langle F \rangle_{\Omega}$, and they can be transferred from one compound to another as far as the shape of the basin do not change much.



QTAIM partition in the MgO crystal



Partition of the bulk modulus B and $\kappa = B^{-1}$

- The partition of atomic compressibility is simple:

$$\kappa = \frac{1}{B} = -\frac{1}{V} \left(\frac{\partial V}{\partial p} \right), \quad V = \sum_{\Omega} V_{\Omega}, \quad f_{\Omega} = V_{\Omega}/V,$$

$$\kappa_{\Omega} = \frac{1}{B_{\Omega}} = -\frac{1}{V_{\Omega}} \left(\frac{\partial V_{\Omega}}{\partial p} \right) \Rightarrow \boxed{\kappa = \sum_{\Omega} f_{\Omega} \kappa_{\Omega}}.$$

- This provides a very practical analytical tool. We can see, for instance, that the compressibility of spinels is controlled by the oxide ion (PhysRev B **62** (2000) 13970):


AB_2O_4	B (GPa)	B_A	B_B	B_O	f_O
MgAl ₂ O ₄	215.2	282.1	331.9	201.6	0.8127
MgGa ₂ O ₄	211.2	261.2	283.9	196.1	0.7486
ZnAl ₂ O ₄	214.8	246.0	335.2	203.3	0.7690
ZnGa ₂ O ₄	213.3	241.2	308.6	195.7	0.7070

- $B_O \approx B < B_{Zn} < B_{Mg} < B_{Ga} < B_{Al}$.
- In the static approximation ($T = 0$ K, no ZPE) it *should* be possible to partition all thermodynamic observables into basin contributions.



Influences on B_{Ω} and κ_{Ω} :

- atomic number • polarizability • charge transfer (electronegativity)
- coordination index • crystal structure • what else?



AgGaSe₂ (chalcopyrite phase)

	Ag	Ga	Se
Q	+0.28	+1.00	-0.64


$$c = 31\%, \quad f = 2.6\%$$

Ω	f_{Ω}	κ_{Ω} (TPa ⁻¹)	B_{Ω} (GPa)
Ag	0.238	17.7	56.6
Ga	0.157	22.3	44.8
Se	0.603	19.6	51.1
Total		19.5	51.2

$B(^{31}\text{Ga}) < B(^{34}\text{Se}) < B(^{47}\text{Ag})$ are ordered by atomic number!

We lack a systematic study of the B_{Ω} 's!!!

Elastic equation of state



- For small deformations of the crystal cell around an equilibrium configuration:

$$\phi = \frac{E - E_0}{V_0} = \frac{1}{2} \sum_{ij,kl} c_{ijkl} \epsilon_{ij} \epsilon_{kl} = \frac{1}{2} \sum_{ij,kl} s_{ijkl} \tau_{ij} \tau_{kl}. \quad (1)$$

- Using symmetric deformation components:

$$\epsilon_{ij} = \frac{1}{2} \left[\frac{\partial \delta x_i}{\partial x_j} + \frac{\partial \delta x_j}{\partial x_i} \right], \quad \delta x = x - x_0, \quad x \in \{a, b, c\}. \quad (2)$$

- Strain-stress linear relationships:

$$\tau_{ij} = \frac{\partial \phi}{\partial \epsilon_{ij}}, \quad \tau_{ij} = \sum_{kl} c_{ijkl} \epsilon_{kl}, \quad \epsilon_{ij} = \sum_{kl} s_{ijkl} \tau_{kl}. \quad (3)$$

- Voigt notation:

τ_{11}	τ_{22}	τ_{33}	τ_{23}	τ_{31}	τ_{12}	ϵ_{11}	ϵ_{22}	ϵ_{33}	$2\epsilon_{23}$	$2\epsilon_{31}$	$2\epsilon_{12}$
τ_1	τ_2	τ_3	τ_4	τ_5	τ_6	ϵ_1	ϵ_2	ϵ_3	ϵ_4	ϵ_5	ϵ_6

- Elastic constants and compliances:

$$c_{ij} = \left(\frac{\partial \tau_i}{\partial \epsilon_j} \right)_{\epsilon', 0}, \quad s_{ij} = \left(\frac{\partial \epsilon_i}{\partial \tau_j} \right)_{\tau', 0}, \quad \mathbf{c} = \mathbf{s}^{-1}. \quad (4)$$

How to partition the c_{ij} elastic constants

- The partition of the energy, $E = \sum_{\Omega} E^{\Omega}$, would provide a direct partition of c_{ij} :

$$\phi = \sum_{\Omega} \phi^{\Omega}, \quad \phi^{\Omega} = \frac{E^{\Omega} - E_0^{\Omega}}{V_0} \Rightarrow c_{ij} = \sum_{\Omega} c_{ij}^{\Omega}, \quad c_{ij}^{\Omega} = \left(\frac{\partial^2 \phi^{\Omega}}{\partial \epsilon_i \partial \epsilon_j} \right)_{\epsilon', 0}. \quad (5)$$

However, *we still can't partition the cell energy with the required precision.*

- The indirect way uses the partition of the cell volume:

$$c_{ij} = \left(\frac{\partial \tau_i}{\partial \epsilon_j} \right)_{\epsilon'} = \frac{\partial \tau_i}{\partial V} \frac{\partial V}{\partial \epsilon_j} = \frac{\partial \tau_i}{\partial V} \sum_{\Omega} \frac{\partial V^{\Omega}}{\partial \epsilon_j} = \sum_{\Omega} c_{ij}^{\Omega}, \quad c_{ij}^{\Omega} = \left(\frac{\partial V^{\Omega}}{\partial \epsilon_j} \right) \left(\frac{\partial V}{\partial \tau_i} \right)^{-1} \quad (6)$$


- We need to calculate ($E, V, V^{\Omega} \dots$) for appropriate **finite deformations** of the unit cell: $\mathbf{R} \rightarrow \mathbf{R}' = \mathbf{R}(\mathbf{1} + \epsilon)$. In the case of **cubic systems**:

$$\mathbf{R}' = \begin{pmatrix} a & 0 & 0 \\ 0 & a & 0 \\ 0 & 0 & a \end{pmatrix} \begin{pmatrix} 1 + \epsilon_1 & 2\epsilon_6 & 2\epsilon_5 \\ 2\epsilon_6 & 1 + \epsilon_2 & 2\epsilon_4 \\ 2\epsilon_5 & 2\epsilon_4 & 1 + \epsilon_3 \end{pmatrix} \quad (7)$$


$$V = V_0 \{ 1 + \epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_1 \epsilon_2 + \epsilon_2 \epsilon_3 + \epsilon_3 \epsilon_1 + \epsilon_1 \epsilon_2 \epsilon_3 - 4\epsilon_4^2 - 4\epsilon_5^2 - 4\epsilon_6^2 - 4\epsilon_1 \epsilon_4^2 - 4\epsilon_2 \epsilon_5^2 - 4\epsilon_3 \epsilon_6^2 + 16\epsilon_4 \epsilon_5 \epsilon_6 \} \quad (8)$$




Deformations of the cubic cell



$$\Phi(\epsilon_1) \Rightarrow c_{11}$$






$$\Phi(\epsilon_1, \epsilon_2) \Rightarrow c_{12}, c_{11}$$



$$\Phi(\epsilon_4) \Rightarrow c_{44}$$

$$c_{11} = \sum_{\Omega} c_{11}^{\Omega}, \quad c_{11}^{\Omega} = \left(\frac{\partial V_{\Omega}}{\partial \epsilon_1} \right)_{\epsilon'_1, 0} \left(\frac{\partial V}{\partial \tau_1} \right)^{-1}_{\epsilon'_1, 0} \quad (9)$$




$$V(\epsilon_1, \epsilon'_1 = 0) = V_0(1 + \epsilon_1) \quad (10)$$

		Mg	O	cell
Q_{Ω}	(e)	1,71	-1,71	
V_{Ω}	(bohr ³)	31,70	95,65	127,35
c_{11}^{Ω}	(GPa)	51,50	215,56	267,06



$$c_{12}^{\Omega} = \left(\frac{\partial V_{\Omega}}{\partial \epsilon_2} \right)_{\epsilon', 0} \left(\frac{\partial V}{\partial \tau_1} \right)_{\epsilon', 0}^{-1}, \quad \tau_{\xi} = \left(\frac{\partial \phi}{\partial \epsilon_{\xi}} \right)_{\epsilon', 0}, \quad c_{12}^{\Omega} = \left(\frac{\partial V_{\Omega}}{\partial \epsilon_1} \right)_{\epsilon', 0} \left(\frac{\partial V}{\partial \tau_2} \right)_{\epsilon', 0}^{-1} \quad (11)$$



Ω	c_{12}^{Ω}
Mg	16,87
O	71,13
cell	88,09

$$V(\epsilon_1, \epsilon_2, \epsilon' = 0) = V_0(1 + \epsilon_1 + \epsilon_2 + \epsilon_1 \epsilon_2) \quad (12)$$




Partitioning s_{11} and s_{12}

$$s_{ij} = \left(\frac{\partial \epsilon_i}{\partial \tau_j} \right)_{\tau', 0} \Rightarrow s_{ij} = \sum_{\Omega} s_{ij}^{\Omega}, s_{ij}^{\Omega} = \left(\frac{\partial V^{\Omega}}{\partial \tau_j} \right) \left(\frac{\partial V}{\partial \epsilon_i} \right)^{-1} \bigg|_{\tau', 0} \quad (13)$$

We have tested two different approaches:

- Using the capability of ABINIT for introducing constant stress conditions into the electronic structure calculations.
- Assuming linear stress-strain relationships ($\epsilon = s\tau = c^{-1}\tau$), we determine the cell shape for a given stress condition.

The first approach do not assumes linearity, but both produce equivalent s_{ij} values and partitions.




(TPa ⁻¹)	s_{11}^{Ω}	s_{12}^{Ω}
Mg	0,8628	-0,2132
O	3,6164	-0,8970
Cell	4,4792	-1,1102

Partitioning c_{44} and s_{44}

- $\tau_4 = c_{44}s_4$ and the same $\Phi(\epsilon_4, \epsilon'_4=0)$ data can be used to obtain both c_{44} and s_{44} .
- $\left. \frac{\partial V}{\partial \epsilon_4}, \frac{\partial V^\Omega}{\partial \tau_4} \right|_{\tau'_4,0}$ and $\left. \frac{\partial V}{\partial \tau_4}, \frac{\partial V^\Omega}{\partial \epsilon_4} \right|_{\epsilon'_4,0}$ are **zero** and the formulas previously used give a **0/0 indeterminacy ...**
- ... but we can use L'Hôpital:

$$c_{44} = \sum_{\Omega} \frac{\partial^2 V^\Omega}{\partial \epsilon_4^2} / \left. \frac{\partial^2 V}{\partial \tau_4 \partial \epsilon_4} \right|_{\epsilon'_4,0},$$

$$s_{44} = \sum_{\Omega} \frac{\partial^2 V^\Omega}{\partial \tau_4 \partial \epsilon_4} / \left. \frac{\partial^2 V}{\partial \epsilon_4^2} \right|_{\tau'_4,0}. \quad (14)$$



	c_{44} (GPa)	s_{44} (TPa ⁻¹)
Mg	-25,19	-1,149
O	172,79	+7,915
Cell	147,79	+6,767

We are used to:


- $\mathbf{c}^{-1} = \mathbf{s}$ (General).
- $B = (c_{11} + 2c_{12})/3, \quad \kappa = 3(s_{11} + 2s_{12}), \quad c_{44} = 1/s_{44}$ (Cubic crystals).

However, with the c_{ij}^{Ω} and s_{ij}^{Ω} :


- $c_{ij} = \sum_{\Omega} c_{ij}^{\Omega}$ and $s_{ij} = \sum_{\Omega} s_{ij}^{\Omega}$.
 - $\kappa^{\Omega} = 3(s_{11}^{\Omega} + 2s_{12}^{\Omega}) \dots$
 - $\dots$ but $B^{\Omega} \neq (c_{11}^{\Omega} + 2c_{12}^{\Omega})/3$.
 - $c_{44}^{\Omega} \neq 1/s_{44}^{\Omega}$.
 - $\mathbf{c}^{\Omega} \neq (\mathbf{s}^{\Omega})^{-1}$.




Partitioning in four families of oxide crystals




Perovskite: CaTiO_3



Inverse fluorite: Li_2O



Rock salt: MgO





Cuprite: Cu_2O



Calculation details

- Electronic structure calculations: ABINIT.
- Relativistic norm conserving pseudopotentials (Troulier-Martins).
- The xc functional: PBE96 GGA. PW: $E_{cut} = 41$ hy. $\Delta E \approx 1 \cdot 10^{-5}$ hy.
- Topological analysis: CRITIC2 (qt_{ree}, lev. 8) with interpolations from a $200 \times 200 \times 200$ grid of $\rho(r)$ values.



AMO₃ perovskites

c, f, B	CaTiO ₃			SrTiO ₃			MgSiO ₃		
	65	4,5	162	65	4,9	164	85	11,7	226
Ω	Ca	Ti	$\langle O \rangle$	Sr	Ti	$\langle O \rangle$	Mg	Si	$\langle O \rangle$
V^Ω	89	50	94	111	51	88	38	18	80
Q^Ω	1,63	2,17	-1,27	1,60	2,17	-1,26	1,77	3,27	-1,68
f^Ω	0,21	0,12	0,67	0,26	0,12	0,62	0,13	0,06	0,81
B^Ω	209	206	147	212	208	146	284	196	221
κ^Ω	4,79	4,85	6,80	4,73	4,80	6,86	3,52	5,09	4,52
c_{11}^Ω	52	29	79	65	30	75	35	24	96
c_{12}^Ω	14	8	21	18	8	21	17	12	46
c_{44}^Ω	23	-29	47	25	-30	33	37	-65	68

Units: c, f (%); V (bohr³); Q (e); B, c_{ij} (GPa); κ, s_{ij} (TPa⁻¹).



Conclusions

- *The QTAIM partitioning of the c_{ij} and s_{ij} elastic constants is possible by following the evolution of the cell energy and volume, E and V , and that of the basin volumes, V^Ω , under appropriate finite deformations of the unit cell.*
- *Keeping the stresses constant to get the s_{ij} compliances is possible either by using the linear strain-stress relationship ($\epsilon = s\tau$) or by imposing constant stress conditions in the electronic structure calculations, like the ABINIT code does.*
- *A good precision on the basin volumes is required, particularly in the computation of the delicate shear moduli. This has been possible with the new techniques developed for CRITIC2.*
- *The s_{ij}^Ω fulfill the internal consistency rules. For instance: $\kappa_\Omega = 3(s_{11} + 2s_{12})$.*
- *We are still working on the rules that govern the basin partition.*
- *Are you interested in any important case?*



- Funds have been provided by the Spanish Ministerio de Ciencia e Innovación under grant CTQ2009-08376.
- The research is part of the MALTA Consolider Ingenio initiative (grant no. CSD2007-00045). See <http://www.malta-consolider.com/>.





Dedicated to the memory of Miguel Álvarez Blanco (1969–2010), who left behind two small children and a beloved wife.

